



# The Methionine Synthase Reductase (MTRR) A66G and Methionine Synthase (MTR) A2756G Polymorphisms Might be a Risk Factor for Colorectal Cancer in People Living near the Southern Coast of the Caspian Sea

## ARTICLE INFO

### Article Type

Original Research

### Authors

Farkhondeh Nemati<sup>1\*</sup>  
 Fatemeh Malakpour<sup>2</sup>  
 Rohullah Abdollahpour<sup>3</sup>  
 Mahboubeh Rajabzadeh<sup>4</sup>  
 Maryam Gholampour<sup>5</sup>

1. Assistant Professor in Animal Physiology, Department of Biology, Faculty of Basic Science, Islamic Azad University, Qaemshahr Branch, Mazandaran, Iran
2. M.Sc. Student of Biology, Islamic Azad University, Qaemshahr Branch, Mazandaran, Iran..
3. Assistant Professor in Animal Genetics and Breeding, Department of Animal Science, Islamic Azad University, Qaemshahr Branch, Mazandaran, Iran.
4. M.Sc. Student of Biology, Islamic Azad University, Qaemshahr Branch, Mazandaran, Iran.
5. M.Sc. Student of Control Engineering, K. N. Toosi University of Technology, Department of Electrical Engineering, Tehran, Iran.

### \*Corresponding authors:

Assistant Professor in Animal Physiology,  
 Department of Biology, Faculty of Basic  
 Science, Islamic Azad  
 University, Qaemshahr Branch,  
 Mazandaran, Iran,  
 Farkhondehnemati@gmail.com

### Article History

Received: 2021/09/22

Accepted: 2022/01/12

ePublished: March 6, 2021

## ABSTRACT

A case-control study was designed to investigate the association between the risk of colorectal cancer and genetic variation in four single-nucleotide polymorphism (SNPs) within involved genes in folate metabolism: C677T and A1298C of methylenetetrahydrofolate reductase (MTHFR), A66G of methionine synthase reductase (MTRR), and A2756G of methionine synthase (MTR). Genomic DNA was extracted from peripheral blood of 50 patients with newly diagnosed colorectal cancer and 100 non-cancer controls. The statistical analysis was carried out by logistic regression. The results showed that the MTR and MTRR SNPs were significantly associated with increased risk of colorectal cancer ( $p < 0.01$ ). Moreover, no significant relation was found between MTHFR C677T and MTHFR A1298C polymorphisms and the risk of colorectal cancer. These findings suggest that the MTRR A66G and MTR A2756G polymorphisms might be some genetic risks factor for colorectal cancer in the studied population.

**Keywords:** Colorectal Cancer; Gene Polymorphism; Methylenetetrahydrofolate Reductase; Methionine Synthase Reductase; Methionine Synthase


Copyright© 2020, TMU Press. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms

## 1. Introduction

Folate has shown a key role in the etiology of colorectal cancer [1]. The leading cause of colorectal cancer might be the genetic differences in the folate metabolic pathways, such as single nucleotide polymorphisms (SNPs) [2]. Folate converts into one of the three forms within the cells, namely 5-methyl-THF (Tetrahydrofolate), 5,10 methylene-THF, or 10-formyl-THF. In this regard, 5-methyl-THF is the dominant circulating form of folate, which is used for processes such as DNA methylation reactions. Also, 5-methyl-THF donates a methyl group to homocysteine which is converted it to methionine. Methionine is later catabolized to produce S-adenosylmethionine (SAM), a major source of methyl groups used for DNA methylation. The second folate form (i.e., 5,10 methylene-THF) changes deoxyuridylate (dUMP) into thymidylate (dTMP) and the last folate form (i.e., 10-formyl-THF) produces new purines. Purines are used both in RNA and DNA synthesis. Furthermore, dTMP is produced for DNA synthesis and DNA repair or DNA replication [3] (Figure 1). The enzyme that converts 5,10 methylene-THF to 5-methyl-THF is MTHFR (Figure 1). The MTHFR gene includes two polymorphisms: C677T and A1298C. The MTHFR variant C677T causes a decrement in enzyme activity, leading to differential methylation of genomic DNA [4,5], which is associated with different forms of cancer. Also, the incremental change in the plasma level of homocysteine could be caused by reduced MTHFR activity. Homocysteine may also affect methylation of DNA [6]. The second MTHFR variant (i.e., A1298C) also causes conformational modifications within the MTHFR enzyme that alter the enzymatic activity [7,8]. The remethylation of homocysteine to methionine which is needed for SAM production is catalyzed by MTR (A66G). In methylation process, vitamin B<sub>12</sub> is used as a cofactor (A66G) by MTR. MTRR (A2756G) may catalyze the reductive methylation of vitamin B<sub>12</sub> using SAM as a methyl donor and leads to the activation of MTR [9,10].

The deficiency of folate metabolism due to dietary, pharmacological, and genetic factors has been suggested to increase the risk of colorectal cancer via aberrations in DNA methylation and

imbalance of DNA precursors [11,12]. In this respect, some environmental factors such as geographical and ethnic backgrounds are extremely important to capture biodiversity and fully understanding risk and treatments related to polymorphism [13]. In the present study, we proposed to examine the correlations between polymorphism of C677T and A1298C within methylenetetrahydrofolate reductase (MTHFR), A66G within methionine synthase reductase (MTRR), and A2756G within methionine synthase (MTR) and colorectal cancer occurrence in people living in the South of Caspian Sea (Mazandaran province, Iran).



**Figure 1.** Folate cycle and the role of methylenetetrahydrofolate reductase (MTHFR), methionine synthase reductase (MTRR), and methionine synthase (MTR) gene in the metabolism of folate (DHF, dihydro tetra folic acid; THF, tetrahydrofolic acid; 5-10-MethylenTHF, 5-10-methylenetetrahydrofolate; 5-Methyl THF, 5-methyltetrahydrofolate; SAM, S-adenosylmethionine).

## 2. Materials and Methods

### 2.1. Study population and sample collection

50 patients with nonhereditary colorectal cancer diagnosed after a colonoscopy at Imam Hospital (Sari, Iran) between April 2019 and August 2020 were study cases of this experiment. The controls included 100 blood donors with the same age and blood collection year. Each patient gave us their Informed consent. The study was approved by the Medical Ethics Committee of Sari Medical Sciences Branch, Islamic Azad University. The mean age of the patients was 55±18 years for women and 59±15 years for men. People with a history of various risk factors for chronic disease

and other types of genetic and familial malignancies were excluded. All samples were collected randomly. Blood samples were collected in EDTA tubes and later were immediately centrifuged and separated for plasma and stored at -20°C. The DNA was extracted from whole blood samples by DNG plus Genomic DNA extraction kit (Cinnagen, Iran) according to the manufacturer's instructions..

## 2.2. Genotyping of the MTHFR, MTR, and MTRR genes

Genotyping for four SNPs (MTHFR C677T, MTHFR A1298C, MTR A2756G, and MTRR A66G) was carried out using polymerase chain reaction (PCR) in order to amplify blood DNA followed by restriction enzyme digestion and gel electrophoresis. DNA amplification of C677T fragment by PCR was performed using two primers: The forward primer and the reverse primer. The fragment was amplified in 25 µl PCR mixture containing 1 µM of template DNA, 0.2 µl *Tag* DNA polymerase, 0.5 µl dNTPs mix, 1 µM forward primer, 1 µM reverse primer, and 1.5 µM MgCl<sub>2</sub>. Polymorphisms were identified by the enzymatic digestion of the obtained fragment by

PCR amplification with a restriction endonuclease. The polymorphisms create a restriction site for the enzymes. The enzymatic digestion, which was acquired in a 10-µl reaction mixture, consisted of the following materials: 1X NE buffer (50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, 1 mM DTT, pH 7.9@ 25°C) and 5 units of *HinfI*/ 10µl reaction mixture. The mixture was incubated at 37°C for enzymatic digestion for 3 h. The products of enzymatic digestion were separated by electrophoresis in 3% agarose gel stained with ethidium bromide. The sequences of the primers, annealing temperatures, restriction enzymes, genotype, and length of the restriction fragments (bp) are listed in Table 2.

**Table 1.** Background Characteristics of Patients by Age and Sex

| Characteristic | n (%)   |
|----------------|---------|
| Total patients |         |
| Males          | 28(56)  |
| Females        | 22(44)  |
| Age(in years)  |         |
| Mean           | 57±16.2 |
| Median         | 57      |
| Range          | 37-74   |

**Table 2.** The Sequences of the Primers, Annealing Temperatures (T<sub>m</sub>), Restriction Enzymes, Genotype, and Length of the Restriction Fragments (bp)

| SNPs         | Primer sequences                                                           | T <sub>m</sub> | Restriction enzymes | Genotype and Fragment size (bp) |                 |            |
|--------------|----------------------------------------------------------------------------|----------------|---------------------|---------------------------------|-----------------|------------|
|              |                                                                            |                |                     | Normal Homozygous               | Heterozygous    | Homozygous |
| MTHFR C677T  | F:(5'-TGAAGGAGAAGGTGTCTGCGGA-3')<br>R:(5'-AGGACGGTGCCTGAGAGTG-3')          | 59°C           | <i>HinfI</i>        | CC(86 bp)                       | CT(86,50,36)    | TT(50, 36) |
| MTHFR A1298C | F:(5'-GAGCTGAAGGACTACTACCTC-3')<br>R:(5'-CACTTTGTGACCATTCGGTTG-3')         | 61°C           | <i>MboII</i>        | AA(25,28,31)                    | AC(59,31,28,25) | CC(59,25)  |
| MTR A2756G   | F:(5'-GAAGTAGAAGACAGAAATTCTCTA-3')<br>R:(5'-CATGGAAGAATATCAAGATATTAGA-3')  | 61°C           | <i>HaeIII</i>       | AA(66)                          | AG(66,37,29)    | GG(37,29)  |
| MTRR A66G    | F:(5'-CAGGCAAAGGCCATCGCAGAAGACAT-3')<br>R:(5'-CACTTCCAACCAAAATCTTCAAAG-3') | 59°C           | <i>NdeI</i>         | AA(44,22)                       | AG(66,44,22)    | GG(66)     |

T<sub>m</sub>: Annealing temperature; F: Forward; R: Reverse; bp: base pairs

## 2.3. Statistical analysis

Statistical analyses and computations were done using the R software [14]. Allele and genotype frequencies and Hardy-Weinberg equilibrium for the SNPs were estimated using the GENETICS package [15] in R. Associations between SNPs

genotypes and colorectal cancer were performed using a multivariable logistic regression model including all four SNPs genotypes, sex and age variables to adjust their confounding effects. Odds ratios and their 95% confidence interval were calculated based on the output of logistic regression analysis. All p-values less than 0.05 were

considered as statistically significant.

## 2.4. Ethical statement

This study was approved by the Institution Ethics Committee approval with the number of IR.IAU.SARI.REC.1398.079 and all participants provided informed consent. The online version of the approved version is available to the public at the address <http://ethics.research.ac.ir/IR.IAU.SARI.REC.1398.079>.

## 3. Results

### 3.1. Distribution of genotypes in patients and controls

The general characteristics of the studied patients are presented in Table 1. The frequencies of alleles and genotypes of the MTHFR C677T, MTHFR A1298C, MTR A2756, and MTRR A66G loci are shown in Table 3. The chi-square ( $\chi^2$ ) test demonstrated that the distributions of MTRR and MTHFR A1298C loci genotypes deviated from Hardy-Weinberg equilibrium ( $p < 0.05$ ). Association parameters (odds ratios (OR), 95% confidence intervals (CI), and p-values) adjusted for the effects of age and sex are given in Table 3. We examined the relation between the four common polymorphisms of three critical enzymes involved in folate metabolism and the risk of colorectal cancer. Our result suggested that no significant relation was found between MTHFR C677T and MTHFR A1298C polymorphism and the risk of colorectal cancer. The presence of C677T mutation (CT-genotype, TT-genotype and T-allele) was not a significant risk factor for colorectal cancer. Results for patients with the A1298C mutation (AC-genotype, CC-genotype and C-allele) is similar to the C677T mutation. Unlike MTHFR C677T and MTHFR A1298C, the genotype and allele frequencies for MTR A2756G and MTRR A66G differed significantly between patient and control groups. In MTR A2756G polymorphism, GG genotype had the highest risk (OR = 7.41, 95%CI = 1.96-27.9,  $p = 0.003$ ) for the colorectal cancer. Genotypes carrying the G allele (AG + GG) also showed significant risk (OR = 3.18, 95%CI = 1.34-7.52,  $p = 0.008$ ). The G allele was significantly associated with colorectal cancer (OR = 2.55, 95%CI = 1.36-4.80,  $p = 0.003$ ). Moreover, in MTRR A66G polymorphism, the GG genotype was associated with an increased risk of colorectal cancer (OR = 107.6, 95%CI = 12.45-929.9,  $p < 0.0001$ ). The carriers of G (AG + GG) were at a high

risk (OR = 9.34, 95%CI = 4.06-21.50,  $p < 0.0001$ ). Finally, a strong association was found between allele G and colorectal cancer (OR = 6.63, 95%CI = 3.40-12.94,  $p < 0.0001$ ).

**Table 3.** Genotype Frequencies in Case and Control Groups, Odds Ratios (OR) and 95% CIs for MTRR, MTR, and MTHFR Polymorphisms Adjusted for the Effects of Sex and Age

| Genotype/Allele | Control<br>n=100<br>(%) | Case<br>n=50<br>(%) | OR    | 95%CI       | P<br>value  |
|-----------------|-------------------------|---------------------|-------|-------------|-------------|
| MTHFR<br>C677T  |                         |                     |       |             | 0.39        |
| CC              | 69 (69)                 | 32 (64)             | 1     | Reference   | -           |
| CT              | 30 (30)                 | 16 (32)             | 1.11  | 0.40-3.07   | 0.84        |
| TT              | 1 (1)                   | 2 (4)               | 3.49  | 0.17-70.6   | 0.41        |
| CT + TT         | 31(31)                  | 18(36)              | 1.09  | 0.44-2.71   | 0.85        |
| C               | 168(84)                 | 80(80)              | 1     | Reference   | -           |
| T               | 32(16)                  | 20(20)              | 1.22  | 0.49-3.04   | 0.66        |
| MTHFR<br>A1298C |                         |                     |       |             | 0.57        |
| AA              | 81 (81)                 | 34 (68)             | 1     | Reference   | -           |
| AC              | 16 (16)                 | 12 (24)             | 1.65  | 0.51-5.30   | 0.40        |
| CC              | 3 (3)                   | 4 (8)               | 2.31  | 0.32-16.4   | 0.40        |
| AC + CC         | 19(19)                  | 16(32)              | 1.97  | 0.74-5.26   | 0.17        |
| A               | 178(89)                 | 80(8)               | 1     | Reference   | -           |
| C               | 22(11)                  | 20(2)               | 1.54  | 0.68-3.49   | 0.29        |
| MTR<br>A2756G   |                         |                     |       |             | 0.007       |
| AA              | 60 (60)                 | 17 (34)             | 1     | Reference   | -           |
| AG              | 34 (34)                 | 21 (42)             | 2.72  | 1.01-7.40   | 0.049       |
| GG              | 6 (6)                   | 12 (24)             | 7.41  | 1.96-27.9   | 0.003       |
| AG + GG         | 40(40)                  | 44(66)              | 3.18  | 1.34-7.52   | 0.008       |
| A               | 154(77)                 | 55(55)              | 1     | Reference   | -           |
| G               | 46(23)                  | 45(45)              | 2.55  | 1.36-4.80   | 0.003       |
| MTRR<br>A66G    |                         |                     |       |             | <0.00<br>01 |
| AA              | 83 (83)                 | 17 (34)             | 1     | Reference   | -           |
| AG              | 16 (16)                 | 16 (32)             | 3.78  | 1.44-9.88   | 0.007       |
| GG              | 1 (1)                   | 17 (34)             | 107.6 | 12.45-929.9 | <0.00<br>01 |
| AG + GG         | 17(17)                  | 33(66)              | 9.34  | 4.06-21.50  | <0.00<br>01 |
| A               | 182(91)                 | 50(50)              | 1     | Reference   | -           |
| G               | 18(9)                   | 50(50)              | 6.63  | 3.40-12.94  | <0.00<br>01 |

## 4. Discussion

Colorectal cancer rates third among the most common cancers with 1.8 million new cases and the fourth rate in the most common causes of

cancer death in 2018 in the world [16]. According to data from the National Cancer Registry of Iran and the Center for Disease Control and Prevention of the Ministry of Health and Medical Education of Iran, the central, northern, and western provinces of Iran have the highest incidence rates of colorectal cancer [17]. However, it is the most often diagnosed cancer and the main cause of cancer death. The mortality rate of the disease varies across the countries and depends on the degree of economic development, social and lifestyle factors [16]. In addition to environmental factors, genetic factors such as genetic polymorphisms of enzymes involved in folate metabolism might be the risk factors for this cancer. Among these enzymes, MTHFR, MTR, and MTRR are risky genes.

In the present study, the associations of MTHFR C677T and A1298C, MTRR A66G, and MTR A2756G polymorphisms with the susceptibility of colorectal cancer were analyzed. The results showed that the MTR 2756AG genotype, 2756GG genotype, 2756 AG and GG genotype, and 2756 G-allele were significantly associated with increased risk of colorectal cancer and MTRR 66AG genotype, 66GG genotype, 66 AG and GG genotype, and 66 G-allele are strong risk factors for colorectal cancer. No significant association was found between MTHFR C677T and MTHFR A1298C polymorphisms and the risk of colorectal cancer.

There are conflicting results from previous colorectal cancer studies about the relationship between the polymorphisms of the three investigated genes and the risk of colorectal cancer. Several studies have shown the associations between genotypes of MTHFR, MTRR, and MTR with increased risk of colorectal cancer [18-22] but in other studies no relationship was found [23]. On the contrary, some studies have reported even reverse associations with reduced risk of colorectal cancer and even protective effect on it [24-30] depending on adequate dietary folate intake. Furthermore, some studies show low plasma folate levels were associated with a lower risk of colorectal cancer [31]. They concluded that folic acid supplementation (1 mg/d) did not prevent the recurrence of colorectal adenomas and tended to increase greater adenoma multiplicity.

Nevertheless, in another trial, folic acid supplementation was associated with lower recurrence of adenomas among those with low plasma folate at baseline [32]. In a study of nutrients, foods, and colorectal cancer results manifested that folate intake lasted through diet or supplements had a complex relationship with colorectal cancer risk; it potentially promoted the growth of pre-existing tumors while inhibited the formation of new tumors in healthy tissue [33]. Mokarram *et al.* showed that the MTHFR 677TT genotype combined with a high serum folate status might be a risk factor for tumor-specific gene promoter hypermethylation among colorectal patients of southern Iran [34].

In another study by Naghibalhossaini *et al.* on the same people, MTHFR 667T allele was strongly correlated with colorectal cancer and microsatellite instability. They proposed that it might be mediated by the inactivation of mismatch repair genes caused by their promoter hypermethylation. They also found no notable difference in the MTHFR 1298C allele. They concluded that it could be due to no significant alteration in the activity of the enzyme caused by this polymorphism [35].

Similar to multifactor disorders, colorectal cancer is affected by both genetic and environmental factors and the effect of genotype may differ in populations with different amounts of folate intake [36]. So, many factors, like dietary folate supply, alcohol intake, age, and gender may modify the effects of these gene polymorphisms and the risk of colorectal cancer [37].

polymorphisms measurement in four SNPs of three genes simultaneously was the aim of in this case-control study, however, the present study suffered from some limitations; e.g., the lack of measurement of plasma folate and folate levels in red blood cells and folate in the diet. Another limitation of the present research was the few number of examined samples due to executive constraints. The few examining population could reduce the power of the statistical analysis. Therefore, to avoid further barrier, we did not consider further analyses in each subgroup. Since this study is not generalized, the results should be explicated cautiously. Hence, it is recommended to perform similar studies with an extended sample size and further evaluation in terms of

micronutrient status for various ethnic groups.

**Declaration of interest: none**

### Acknowledgments

This work was supported in part by a grant from Islamic Azad University, Qaemshahr Branch, with grant number 1.17075.

### References

- [1] Macrae FA. Colorectal cancer: Epidemiology, risk factors, and protective factors. Uptodate com [ažurirano 9. lipnja 2017. 2016 Jan.
- [2] Pabalan, N., Singian, E., Tabangay, L., Jarjanazi, H. and Singh, N., 2015. Associations of the A66G Methionine Synthase Reductase Polymorphism in Colorectal Cancer: A Systematic Review and Meta-Analysis: Supplementary Issue: Biomarkers for Colon Cancer. *Biomarkers in cancer*, 7, pp.BIC-S25251.
- [3] Scotti M, Stella L, Shearer EJ, Stover PJ. Modeling cellular compartmentation in one-carbon metabolism. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*. 2013 May;5(3):343-65. doi: 10.1002/wsbm.1209.
- [4] Yadav, S., Longkumer, I., Joshi, S. and Saraswathy, K.N., 2021. Methylenetetrahydrofolate reductase gene polymorphism, global DNA methylation and blood pressure: a population based study from North India. *BMC Medical Genomics*, 14(1), pp.1-8.
- [5] Crider, K.S., Yang, T.P., Berry, R.J. and Bailey, L.B., 2012. Folate and DNA methylation: a review of molecular mechanisms and the evidence for folate's role. *Advances in nutrition*, 3(1), pp.21-38.
- [6] Barroso, M., Handy, D.E. and Castro, R., 2019. The link between hyperhomocysteinemia and hypomethylation: Implications for cardiovascular disease. *Journal of Inborn Errors of Metabolism and Screening*, 5.
- [7] Wan, L., Li, Y., Zhang, Z., Sun, Z., He, Y. and Li, R., 2018. Methylenetetrahydrofolate reductase and psychiatric diseases. *Translational psychiatry*, 8(1), pp.1-12.
- [8] Petrone, I., Bernardo, P.S., Dos Santos, E.C. and Abdelhay, E., 2021. MTHFR C677T and A1298C Polymorphisms in Breast Cancer, Gliomas and Gastric Cancer: A Review. *Genes*, 12(4), p.587.
- [9] Kim, J.W., Jeon, Y.J., Jang, M.J., Kim, J.O., Chong, S.Y., Ko, K.H., Hwang, S.G., Oh, D., Oh, J. and Kim, N.K., 2015. Association between folate metabolism-related polymorphisms and colorectal cancer risk. *Molecular and clinical oncology*, 3(3), pp.639-648.
- [10] Lyon, P., Strippoli, V., Fang, B. and Cimmino, L., 2020. B vitamins and one-carbon metabolism: implications in human health and disease. *Nutrients*, 12(9), p.2867.
- [11] Guraya, S.Y., 2014. Chemopreventive role of vitamin D in colorectal carcinoma. *Journal of Microscopy and Ultrastructure*, 2(1), pp.1-6.
- [12] Nolfo, F., Rametta, S., Marventano, S., Grosso, G., Mistretta, A., Drago, F., Gangi, S., Basile, F. and Biondi, A., 2013. Pharmacological and dietary prevention for colorectal cancer. *BMC surgery*, 13(2), pp.1-11.
- [13] Izmirli M. A literature review of MTHFR (C677T and A1298C polymorphisms) and cancer risk. *Molecular biology reports*. 2013 Jan 1;40(1):625-37. doi: 10.1007/s11033-012-2101-2.
- [14] Team RC. R: A language and environment for statistical computing.
- [15] Warnes G, Gajabara G, Leisch F, Man M. genetics: Population Genetics. R package version 1.3. 8.1. The Comprehensive R Archive Network. 2013.
- [16] Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*. 2018 Nov;68(6):394-424. doi: 10.3322/caac.21492.
- [17] Shadmani FK, Ayubi E, Khazaei S, et al. Geographic distribution of the incidence of colorectal cancer in Iran: a population-based study. *Epidemiology and health*. 2017;39. Doi: 104178/epih.e2017020
- [18] Ghorbani, M., Azghandi, M., Khayami, R., Baharara, J. and Kerachian, M.A., 2021. Association of MTHFR C677T variant genotype with serum folate and Vit B12 in Iranian patients with colorectal cancer or adenomatous polyps. *BMC Medical Genomics*, 14(1), pp.1-8.
- [19] Yari, A., Meybodi, S.M.E., Karam, Z.M., Samoudi, A., Hashemi, F., Aalipour, M., Abadi, M.F.S. and Dabiri, S., 2021. Association of MTHFR 677C> T and 1298A> C genetic polymorphisms with colorectal cancer: genotype and haplotype analysis in a Southeast Iranian population. *Gene Reports*, p.101399.
- [20] Panprathip, P., Petmitr, S., Tungtrongchitr, R., Kaewkungwal, J. and Kwanbunjan, K., 2019. Low folate status, and MTHFR 677C> T and MTR 2756A> G polymorphisms associated with colorectal cancer risk in Thais: a case-control study. *Nutrition Research*, 72, pp.80-91.

- [21] Wettergren, Y., Odin, E., Carlsson, G. and Gustavsson, B., 2010. MTHFR, MTR, and MTRR polymorphisms in relation to p16INK4A hypermethylation in mucosa of patients with colorectal cancer. *Molecular medicine*, 16(9), pp.425-432.
- [22] Xu, L., Qin, Z., Wang, F., Si, S., Li, L., Lin, P., Han, X., Cai, X., Yang, H. and Gu, Y., 2017. Methylenetetrahydrofolate reductase C677T polymorphism and colorectal cancer susceptibility: a meta-analysis. *Bioscience reports*, 37(6).
- [23] Theodoratou E, Farrington SM, Tenesa A, et al. Dietary vitamin B6 intake and the risk of colorectal cancer. *Cancer Epidemiology and Prevention Biomarkers*. 2008 Jan 1;17(1):171-82. doi: 10.1158/1055-9965.EPI-07-0621.
- [24] Zhou D, Mei Q, Luo H, Tang B, Yu P. The polymorphisms in methylenetetrahydrofolate reductase, methionine synthase, methionine synthase reductase, and the risk of colorectal cancer. *International journal of biological sciences*. 2012;8(6):819. doi: 10.7150/ijbs.4462.
- [25] Cui LH, Shin MH, Kim HN, et al. Methylenetetrahydrofolate reductase C677T polymorphism in patients with lung cancer in a Korean population. *BMC medical genetics*. 2011 Dec 1;12(1):28. doi: 10.1186/1471-2350-12-28.
- [26] Pande, M., Chen, J., Amos, C.I., Lynch, P.M., Broaddus, R. and Frazier, M.L., 2007. Influence of methylenetetrahydrofolate reductase gene polymorphisms C677T and A1298C on age-associated risk for colorectal cancer in a caucasian lynch syndrome population. *Cancer Epidemiology and Prevention Biomarkers*, 16(9), pp.1753-1759.
- [27] Huang, Y., Han, S., Li, Y., Mao, Y. and Xie, Y., 2007. Different roles of MTHFR C677T and A1298C polymorphisms in colorectal adenoma and colorectal cancer: a meta-analysis. *Journal of human genetics*, 52(1), pp.73-85.
- [28] Pardini B, Kumar R, Naccarati A, et al. MTHFR and MTRR genotype and haplotype analysis and colorectal cancer susceptibility in a case-control study from the Czech Republic. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*. 2011 Mar 18;721(1):74-80. doi:10.1016/j.mrgentox.2010.12.008.
- [29] Fernández-Peralta AM, Daimiel L, Nejdá N, et al. Association of polymorphisms MTHFR C677T and A1298C with risk of colorectal cancer, genetic and epigenetic characteristic of tumors, and response to chemotherapy. *International journal of colorectal disease*. 2010 Feb 1;25(2):141-51. doi: 10.1007/s00384-009-0779-y.
- [30] Shiao, S.P.K., Lie, A. and Yu, C.H., 2018. Meta-analysis of homocysteine-related factors on the risk of colorectal cancer. *Oncotarget*, 9(39), p.25681.
- [31] Lee JE, Wei EK, Fuchs CS, et al. Plasma folate, methylenetetrahydrofolate reductase (MTHFR), and colorectal cancer risk in three large nested case-control studies. *Cancer, Causes Control*, Pap. Symp. 2012 Apr 1;23(4):537-45. doi: 10.1007/s10552-012-9911-3.
- [32] Guerreiro CS, Carmona B, Gonçalves S, et al. Risk of colorectal cancer associated with the C677T polymorphism in 5, 10-methylenetetrahydrofolate reductase in Portuguese patients depends on the intake of methyl-donor nutrients. *The American journal of clinical nutrition*. 2008 Nov 1;88(5):1413-8. doi: 10.3945/ajcn.2008.25877.
- [33] Song M, Garrett WS, Chan AT. Nutrients, foods, and colorectal cancer prevention. *Gastroenterology*. 2015 May 1;148(6):1244-60. doi: 10.1053/j.gastro.12.035.
- [34] Mokarram P, Naghibalhossaini F, Firooz MS, et al. Methylenetetrahydrofolate reductase C677T genotype affects promoter methylation of tumor-specific genes in sporadic colorectal cancer through an interaction with folate/vitamin B12 status. *World J. Gastroenterol: WJG*. 2008 Jun 21;14(23):3662. doi: 10.3748/wjg.14.3662.
- [35] Naghibalhossaini F, Mokarram P, Khalili I, et al. MTHFR C677T and A1298C variant genotypes and the risk of microsatellite instability among Iranian colorectal cancer patients. *Cancer genetics and cytogenetics*. 2010 Mar 1;197(2):142-51. doi: 10.1016/j.cancergencyto.2009.11.014.
- [36] Kim DH. The interactive effect of methyl-group diet and polymorphism of methylenetetrahydrofolate reductase on the risk of colorectal cancer. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*. 2007 Sep 1;622(1-2):14-8. doi:10.1016/j.mrfmmm.2007.04.012.
- [37] Lima CS, Nascimento H, Bonadia LC, et al. Polymorphisms in methylenetetrahydrofolate reductase gene (MTHFR) and the age of onset of sporadic colorectal adenocarcinoma. *International journal of colorectal disease*. 2007 Jul 1;22(7):757. doi: 10.1007/s00384-006-0237-z.