

# **Reviewing the effect of Acute Sleep Deprivation on Functional Connectivity of Reward Network and Adverse Emotional Experience**

### **ARTICLE INFO**

Article Type: Original Paper

Authors: Hanieh Riazi<sup>1\*</sup>

1. Department of Physiology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran.

\* *Corresponding author:* Hanieh Riazi

E-mail: hanieh.riazi@modares.ac.ir

## ABSTRACT

While the precise functions of sleep are not fully understood, it is an important part of daily life and involves a series of events that follow a consistent night-time cycle and enable the human body to function at its best. More than 30% of adults suffer from sleep deprivation (SD). SD can lead to negative effects on cognitive function, including reward and emotional processing. Here, we review the consequences of acute SD on reward network and emotional processing, and the activity and connectivity of different brain regions involved in reward network and emotional processing, with a focus on neuroimaging studies.

#### **Keywords:**

sleep deprivation, reward network, emotional processing, brain networks, functional connectivity, neuroimaging.

Copyright© 2020, TMU Press. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms.

#### Introduction

Sleep is a vital physiological process that plays an essential role in maintaining both physical and mental health [1]. It is marked by a significant reduction in consciousness and responsiveness to sensory stimuli, both of which can be rapidly reversible [2]. Sleep affects almost every bodily and mental function and causes intricate alterations in the physiology of numerous systems [3]. The body performs the necessary processes to support brain function and general health while you sleep [3]. At various stages of sleep, important processes like neurogenesis and synaptogenesis take place [4]. Sleep is also essential for removing toxins that build up around brain cells while we are awake [4]. Cerebrospinal fluid production and circulation rise significantly while you sleep, aiding in the removal of toxins like beta-amyloid from brain tissues [4]. Sleep is also essential for preserving mental balance and controlling emotions [4]. Getting enough sleep is essential for all cognitive processes, including consolidation, learning, memory attention. language, reasoning, creativity, and decisionmaking [4].

Brain activity undergoes significant changes during both non-rapid eye movement and rapid eye movement sleep stages [5]. Sleep is therefore anticipated to be essential for brain function [6]. Older adults usually need 8 to 9 h of sleep per night, while healthy adults need 7 to 9 h [1]. The most prevalent sleep-related complaint, sleep deprivation (SD), can result from losing even one hour of sleep [7]. A reduction in the typical length or quality of sleep is a hallmark of SD [7]. If SD lasts for one or two days, it is classified as acute SD, while prolonged deprivation is referred to as chronic SD [8]. The severity of SD can differ among individuals. Acute SD affects more than 30% of the population, and chronic SD is estimated to affect 6-10% globally [8].

People who suffer from SD may suffer severe consequences [9]. In addition to impairing cognitive brain functions like working memory, attention, and decision-making, SD also causes experience to deteriorate [10]. emotional According to studies, SD makes people feel depressed and makes their nerves more sensitive to unpleasant emotional stimuli [11-13]. For instance, increasing waking hours causes agitation, hostility, and anxiety. Furthermore, less control over one's emotions may result in more destructive behavior [10]. Emotional instability can cause major human failure and impair cognitive function, both of which can lead to workplace accidents. Emotional decision-making can be affected by SD. Since rewards and punishments can modify emotions, the emotional state after SD can affect decisions pertaining to gain and loss. Emotional instability following SD and reward network communication, however, has not received much attention in research.

#### Hanieh Riazi –

Additionally, the reward network's alterations and the connection between it and unpleasant emotional experiences are still poorly understood. So here, we review neuroimaging studies investigating how SD causes reduced functional connectivity in the reward network, which is positively associated with negative emotional experiences in humans.

# Effect of acute SD on reward and incentive processing

A network of interrelated brain areas, including the striatum, prefrontal cortex (PFC), hippocampus, amygdala, midbrain ventral tegmental area and nucleus accumbens make up the brain reward system [14]. The striatum receives dopaminergic innervation from the ventral tegmental area [14]. This innervation is linked to and controlled by PFC regions, specifically the inferior orbitofrontal cortex and medial PFC, which direct learning and motivated behavior [14]. It has been repeatedly shown that this system is sensitive to SD, which causes changes in motivated behaviors like impulsivity, risk-taking [14].

By altering dopamine receptor sensitivity and availability in basal ganglia regions, SD impairs dopaminergic function in rats [15, 16]. In a mixed monetary gamble task, humans who have been sleep-deprived for one night exhibit increased ventral striatum activity when they anticipate and receive monetary rewards [17, 12]. Following SD, there is also a significant increase in activity in affect-related frontal cortex regions, such as the insula and medial PFC, that are linked to valuation and viscerosensory functions [12, 17].

These results collectively imply that the state of acute SD appears to hypersensitize subcortical reward-related brain regions as well as associated cortical regions coding salience and valuation [14]. However, there appears to be no difference between rested and deprived conditions in the extent to which functional magnetic resonance imaging (fMRI) signal amplitude tracks reward magnitude in these subcortical and cortical regions [18]. Instead, SD causes a widespread rise in reward sensitivity that undermines the accuracy of reward discrimination, making it harder for the brain to correctly code small increases in reward value from low to high [14].

These findings are supported by the fact that sleep-deprived people's medial PFC, orbitofrontal

cortex, and anterior insula cortex fMRI signals cannot distinguish between trials with monetary reward and punishment values and trials without either [12, 17].

The outcome phase of incentive decision trials also shows this erroneous representation of reward value within frontal regions, indicating a further failure to update accruing reward history and probability [12, 17]. Therefore, after sleep loss, the PFC may not be able to update the changing incentive value of reinforcing stimuli over time due to inaccurate coding of reward and/or punishment valence [12, 17]. This would also lead to less-than-ideal reward-dependent choices and behaviors [14]. According to this profile, people who are sleep-deprived when completing the Iowa Gambling Task take more chances and give more weight to rewards that have been given recently [19, 9]. This pattern, which reflects temporally shorter-sighted reward updating, shows an inability to properly integrate rewards and their increasing value over time after SD [14].

In addition to processing financial rewards, impulsivity is another behavioral trait that is impacted by sleep deprivation [14]. Since impulsivity is a complex concept, there are various ways in which it can show up in behavior [14]. Nonetheless, many incentive-based decision-making processes heavily rely on impulse control [20, 21]. Therefore, after SD, changes in reward decision-making and risktaking may be influenced by changes in impulsivity [14].

Participants who experienced either partial sleep restriction or one night of complete sleep deprivation demonstrated significantly lower response inhibition and cognitive control, more frequent errors of inhibition, and slower learning of cue–incentive associations in tasks that require either cued motor response execution or withholding, also known as a Go/No-Go task [22–26].

According to research to date, SD interferes with reward-value updating and integration and dramatically raises the propensity for reward sensitivity, risk-taking, and impulsivity [27]. Changes in dopamine signaling, which is linked to prolonged wakefulness-possibly more so than sleep deprivation could be one mechanism explaining these increases in approach and consummatory behavior [14]. According to research to date, SD interferes with reward-value updating and integration and dramatically raises the propensity for reward sensitivity, risk-taking, and impulsivity [14]. Changes in dopamine signaling, which is linked to prolonged wakefulness-possibly more so than sleep deprivation could be one mechanism explaining these increases in approach and consummatory behavior [14]. This dopaminergic framework is supported by multiple lines of evidence [14].

First, dopamine is linked to arousal; lower sleep propensity is predicted by naturally higher dopamine levels [28]. Second, research on rodents has shown that wake-promoting stimulant medications, like amphetamine, appear to work in part by preventing dopamine metabolism, which raises dopamine transmission and, consequently, arousal [14]. Third, reducing catecholamines, including dopamine, lowers vigilance, induces sleep, and decreases wake propensity [29, 30].

After SD, changes in reward-driven behavior may also be caused by modifications in dopamine receptors. One night of complete SD has been shown to decrease the availability of dopamine D2 and D3 receptors (D2/3Rs) in the dorsal striatum (which includes the caudate and putamen) as well as the ventral striatum, according to studies using Positron emission tomography ligand [15]. fMRI-measured drops in thalamic activity during an attentional task were predicted by Positron emission tomographymeasured downregulation of D2/3Rs in the ventral striatum [31]. This finding implies that the levels of various dopamine receptors may serve as a proxy for sleep pressure; in line with this, adenosine builds up with longer periods of waking, activating A2A receptors and promoting D2R internalization [15, 32]. Additionally, D2R and A2A receptor agonists exhibit allosteric interaction, meaning that D2R agonists reduce the D2Rs' affinity for their agonists, such as dopamine [33]. Therefore, there are at least two ways that adenosine accumulation, which is linked to longer periods of waking, could reduce activity: D2/3R by increasing D2/3R internalization and by decreasing dopamine binding to D2Rs [14].

Although the downregulation of D2/3Rs associated with SD may seem to contradict the SD-associated increases in neural and behavioral reward sensitivity discussed above, there is a plausible explanation for this paradox involving

D1Rs: D2/3R decreases may result in an imbalance of dopamine receptor availability, which would cause the remaining D1Rs to be disproportionately stimulated by the same amount of presynaptically released dopamine [14]. Outside of SD, this imbalance in rodents leads to increased approach behavior to food rewards [34] and an increased risk of cocaine addiction [35], two behavioral outcomes that have been notably independently linked to SD [36. 371. Additionally, it appears that D1R action is primarily responsible for the rise in striatal fMRI signal brought on by reward incentives under SD [38]. Therefore, conditions SD-associated reductions in D2/3R availability may enhance approach-driven and reward-driven behavior and fMRI-indexed striatal activity by indirectly increasing dopamine binding to the remaining D1Rs [14]. Mechanistic and therapeutic insights into obesity, addiction, and substance use disorders can also be gained from the neural and behavioral effects of sleep deprivation on incentive processes [14].

## Effect of acute SD on emotional processing

In addition to suicidal ideation, suicide attempts, and suicide completion [39–41], sleep deprivation consistently causes changes in negative emotional processing, such as irritability, emotional volatility, anxiety, and aggression [42–45]. These results imply that SD modifies certain process domains, such as basic affective reactivity, as well as more complex ones like emotional expression and discrimination.

The effects of SD on reactivity to simple aversive stimuli were the focus of early neuroimaging studies. For instance, it was demonstrated that a single night of SD increased amygdala reactivity to negative images, including weapons, and poisonous snakes and spiders, by 60% [46]. Notably, rapid, subconscious viewing fear-expressing faces following of sleep restriction has been shown to cause amygdala hyper-reactivity, suggesting that this increased limbic reactivity can happen subconsciously and independently of conscious, deliberate cognition [47].

One possible mechanism underlying such amygdala hypersensitivity involves a loss of regulatory control, resulting in contextually inappropriate amygdala reactivity [48]. Decreases in functional connectivity between top-down

#### Hanieh Riazi -

control regions of the medial prefrontal cortex and the amygdala have been reported following SD [46, 47, 49–51].

The brain's anticipation of cued emotional experiences is further altered by SD [14]. One night of SD, for instance, increases anticipatory responses of the peripheral autonomic nervous system and cue-evoked activity in the amygdala, anterior insula, and anterior cingulate cortex in anticipation of upcoming emotional picture slides [52, 53].

An additional phenotype of the sleep-deprived human brain emerges when participants are challenged with the more complex task of disambiguating between varied emotional signals [14]. Areas of the cingulate, insula, and amygdala linked to salience detection in participants who have had sleep can distinguish between stimuli with varying emotional intensities [14]. In contrast, these areas show a saturated and flattened reaction to emotional stimuli in participants who are sleep deprived [14]. As a result, people who are sleep deprived exhibit a generalized excess of emotional sensitivity along with a reduction in emotional discriminatory specificity [48].

Recent work has begun to uncover the neural basis of these impairments in emotional discrimination. Consistent with the failure to accurately code positive, reward incentive signals. one night of SD impairs the discrimination of faces expressing negative emotions by viscerosensory regions of the anterior insula and anterior cingulate cortices, and to a degree, the subcortical amygdala [54]. Similarly, sleep loss results in generalized, nonspecific increases in amygdala activity in response to aversive and neutral emotional pictures [55, 56].

Due to these results, it has been suggested that SD results in a condition of both central and peripheral emotional hypersensitivity, which makes it impossible to grade emotional reactivity correctly [48]. The afferent-efferent communication between the brain and body, which is essential for the "embodied" mapping of and differentiation between various emotions, is thus compromised by SD [48].

The cellular and molecular processes that underlie the emotional brain and body dysfunction brought on by sleep deprivation are less well understood. The locus coeruleus's elevated noradrenergic tone has been the focus of one theory put forth; sleep deprivation inhibits the typical overnight decrease in noradrenergic tone that typically takes place during REM sleep [48, 57, 58]. Thus, elevated and over-generalized responsiveness may result from the ensuing hypernoradrenergic tone in the amygdala and viscerosensory cortical regions, which are part of the noradrenaline-innervated affective salience network [48]. As a result, affective signaling within this network may lose specificity, which could help to explain the SD-associated inaccuracy in emotional discrimination [48]. It is unlikely, though, that noradrenaline is the only neurochemical component responsible for the affective dysregulation brought on by SD [14]. model that incorporates explanation An sleep-dependent alterations other in neurochemicals, like dopamine, might be more accurate [14].

## Conclusion

In summary, sleep deprivation has a profound effect on how rewards are processed and how emotions are controlled, which drastically changes cognitive and behavioral outcomes. The brain's reward system is impacted by acute sleep deprivation. which impairs judgment and reward sensitivity. Changes increases in dopamine signaling are linked to this altered reward processing, which may lead to an increase in impulsivity and risk-taking behaviors. Anxiety, irritability, and volatility are also increased when sleep deprivation worsens emotional reactivity and impairs the capacity to distinguish between different emotional stimuli. A loss of top-down control and modifications in the functional connectivity between important brain regions, including the amygdala and prefrontal cortex, are the causes of emotional processing dysregulation. All things considered, these results highlight how crucial getting enough sleep is to preserving emotional and cognitive well-being. It is becoming more and more obvious that treatments targeted at increasing the quantity and quality of sleep may be essential for boosting mental health and lowering the risks of emotional dysregulation and poor decision-making as our knowledge of the neurobiological processes underlying the effects of sleep deprivation advances.

# **Conflict of interest**

The author declares no conflicts of interest.

## Acknowledgment

This review article was supported by Medical Faculty of Tarbiat Modares University, the support of this center is appreciated.

## References

- Mason GM, Lokhandwala S, Riggins T, Spencer RMC. Sleep and human cognitive development. Sleep Med Rev 2021; 57:101472.
- 2. Cordi MJ, Rasch B. How robust are sleepmediated memory benefits? Curr Opin Neurobiol 2021; 67:1–7.
- Ferrara M, Gennaro L de. How much sleep do we need? Sleep Med Rev 2001; 5(2):155– 79.
- 4. Riazi H, Nazari M, Raoufy MR, Mirnajafi-Zadeh J, Shojaei A. Olfactory Epithelium Stimulation Using Rhythmic Nasal Air-Puffs Improves the Cognitive Performance of Individuals with Acute Sleep Deprivation. Brain Sci 2024; 14(4).
- Dang-Vu TT, Schabus M, Desseilles M, Sterpenich V, Bonjean M, Maquet P. Functional neuroimaging insights into the physiology of human sleep. Sleep 2010; 33(12):1589–603.
- 6. Orzeł-Gryglewska J. Consequences of sleep deprivation. Int J Occup Med Environ Health 2010; 23(1):95–114.
- 7. Landolt H-P, Holst SC, Sousek A. Effects of acute and chronic sleep deprivation; 2014.
- 8. Vargas I, Nguyen AM, Muench A, Bastien CH, Ellis JG, Perlis ML. Acute and Chronic Insomnia: What Has Time and/or Hyperarousal Got to Do with It? Brain Sci 2020; 10(2).
- Killgore WDS, Balkin TJ, Wesensten NJ. Impaired decision making following 49 h of sleep deprivation. J Sleep Res 2006; 15(1):7– 13.
- Zhang Y, Dai C, Shao Y, Peng J, Yang Y, Hou Y. Decreased Functional Connectivity in the Reward Network and Its Relationship With Negative Emotional Experience After Total Sleep Deprivation. Front Neurol 2021; 12:641810.
- 11. Killgore WD. Sleep Deprivation and

Behavioral Risk-Taking. In: Modulation of Sleep by Obesity, Diabetes, Age, and Diet. Elsevier; 2015. p. 279–87.

- 12. Venkatraman V, Chuah YML, Huettel SA, Chee MWL. Sleep deprivation elevates expectation of gains and attenuates response to losses following risky decisions. Sleep 2007; 30(5):603–9.
- 13. McKenna BS, Dickinson DL, Orff HJ, Drummond SPA. The effects of one night of sleep deprivation on known-risk and ambiguous-risk decisions. J Sleep Res 2007; 16(3):245–52.
- 14. Krause AJ, Simon EB, Mander BA, Greer SM, Saletin JM, Goldstein-Piekarski AN et al. The sleep-deprived human brain. Nat Rev Neurosci 2017; 18(7):404–18.
- 15. Volkow ND, Tomasi D, Wang G-J, Telang F, Fowler JS, Logan J et al. Evidence that sleep deprivation downregulates dopamine D2R in ventral striatum in the human brain. J Neurosci 2012; 32(19):6711–7.
- 16. Tufik S. Changes of response to dopaminergic drugs in rats submitted to REM-sleep deprivation. Psychopharmacology (Berl) 1981; 72(3):257–60.
- Mullin BC, Phillips ML, Siegle GJ, Buysse DJ, Forbes EE, Franzen PL. Sleep deprivation amplifies striatal activation to monetary reward. Psychol Med 2013; 43(10):2215–25.
- Libedinsky C, Smith DV, Teng CS, Namburi P, Chen VW, Huettel SA et al. Sleep deprivation alters valuation signals in the ventromedial prefrontal cortex. Front Behav Neurosci 2011; 5:70.
- 19. Olson EA, Weber M, Rauch SL, Killgore WDS. Daytime Sleepiness Is Associated With Reduced Integration of Temporally Distant Outcomes on the Iowa Gambling Task. Behav Sleep Med 2016; 14(2):200–11.
- 20. Franken IHA, van Strien JW, Nijs I, Muris P. Impulsivity is associated with behavioral decision-making deficits. Psychiatry Res 2008; 158(2):155–63.
- 21. Bechara A, Tranel D, Damasio H. Characterization of the decision-making

deficit of patients with ventromedial prefrontal cortex lesions. Brain 2000; 123 (Pt 11):2189–202.

- 22. Cedernaes J, Brandell J, Ros O, Broman J-E, Hogenkamp PS, Schiöth HB et al. Increased impulsivity in response to food cues after sleep loss in healthy young men. Obesity (Silver Spring) 2014; 22(8):1786–91.
- 23. Demos KE, Hart CN, Sweet LH, Mailloux KA, Trautvetter J, Williams SE et al. Partial sleep deprivation impacts impulsive action but not impulsive decision-making. Physiol Behav 2016; 164(Pt A):214–9.
- 24. Acheson A, Richards JB, Wit H de. Effects of sleep deprivation on impulsive behaviors in men and women. Physiol Behav 2007; 91(5):579–87.
- 25. Ayalon L, Ancoli-Israel S, Drummond SP. Altered brain activation during response inhibition in obstructive sleep apnea. J Sleep Res 2009; 18(2):204–8.
- 26. Anderson C, Platten CR. Sleep deprivation lowers inhibition and enhances impulsivity to negative stimuli. Behav Brain Res 2011; 217(2):463–6.
- 27. Müller SM, Wegmann E, Garcia Arías M, Bernabéu Brotóns E, Marchena Giráldez C, Brand M. Decision Making and Risk Propensity in Individuals with Tendencies towards Specific Internet-Use Disorders. Brain Sci 2022; 12(2).
- 28. Perogamvros L, Schwartz S. The roles of the reward system in sleep and dreaming. Neurosci Biobehav Rev 2012; 36(8):1934– 51.
- 29. McCann UD, Penetar DM, Shaham Y, Thorne DR, Sing HC, Thomas ML et al. Effects of catecholamine depletion on alertness and mood in rested and sleep deprived normal volunteers. Neuropsychopharmacology 1993; 8(4):345– 56.
- Curb JD, Schneider K, Taylor JO, Maxwell M, Shulman N. Antihypertensive drug side effects in the Hypertension Detection and Follow-up Program. Hypertension 1988; 11(3 Pt 2):II51-5.
- 31. Tomasi D, Wang G-J, Volkow ND.

Association between striatal dopamine D2/D3 receptors and brain activation during visual attention: effects of sleep deprivation. Transl Psychiatry 2016; 6(5):e828.

- 32. Elmenhorst D, Meyer PT, Winz OH, Matusch A, Ermert J, Coenen HH et al. Sleep deprivation increases A1 adenosine receptor binding in the human brain: a positron emission tomography study. J Neurosci 2007; 27(9):2410–5.
- 33. Bonaventura J, Navarro G, Casadó-Anguera V, Azdad K, Rea W, Moreno E et al. Allosteric interactions between agonists and antagonists within the adenosine A2A receptor-dopamine D2 receptor heterotetramer. Proc Natl Acad Sci U S A 2015; 112(27):E3609-18.
- 34. Richard JM, Berridge KC. Nucleus accumbens dopamine/glutamate interaction switches modes to generate desire versus dread: D(1) alone for appetitive eating but D(1) and D(2) together for fear. J Neurosci 2011; 31(36):12866–79.
- Park K, Volkow ND, Pan Y, Du C. Chronic cocaine dampens dopamine signaling during cocaine intoxication and unbalances D1 over D2 receptor signaling. J Neurosci 2013; 33(40):15827–36.
- 36. Berro LF, Hollais AW, Patti CL, Fukushiro DF, Mári-Kawamoto E, Talhati F et al. Sleep deprivation impairs the extinction of cocaine-induced environmental conditioning in mice. Pharmacol Biochem Behav 2014; 124:13–8.
- 37. Brondel L, Romer MA, Nougues PM, Touyarou P, Davenne D. Acute partial sleep deprivation increases food intake in healthy men. Am J Clin Nutr 2010; 91(6):1550–9.
- Knutson B, Gibbs SEB. Linking nucleus accumbens dopamine and blood oxygenation. Psychopharmacology (Berl) 2007; 191(3):813–22.
- 39. Bernert RA, Joiner TE. Sleep disturbances and suicide risk: A review of the literature. Neuropsychiatr Dis Treat 2007; 3(6):735–43.
- Kamphuis J, Meerlo P, Koolhaas JM, Lancel M. Poor sleep as a potential causal factor in aggression and violence. Sleep Med 2012; 13(4):327–34.

- 41. Stubbs B, Wu Y-T, Prina AM, Leng Y, Cosco TD. A population study of the association between sleep disturbance and suicidal behaviour in people with mental illness. J Psychiatr Res 2016; 82:149–54.
- 42. Horne JA. Sleep function, with particular reference to sleep deprivation. Ann Clin Res 1985; 17(5):199–208.
- 43. Dinges DF, Pack F, Williams K, Gillen KA, Powell JW, Ott GE et al. Cumulative sleepiness, mood disturbance, and psychomotor vigilance performance decrements during a week of sleep restricted to 4-5 hours per night. Sleep 1997; 20(4):267–77.
- 44. Zohar D, Tzischinsky O, Epstein R, Lavie P. The effects of sleep loss on medical residents' emotional reactions to work events: a cognitive-energy model. Sleep 2005; 28(1):47–54.
- 45. Minkel JD, Banks S, Htaik O, Moreta MC, Jones CW, McGlinchey EL et al. Sleep deprivation and stressors: evidence for elevated negative affect in response to mild stressors when sleep deprived. Emotion 2012; 12(5):1015–20.
- 46. Yoo S-S, Gujar N, Hu P, Jolesz FA, Walker MP. The human emotional brain without sleep--a prefrontal amygdala disconnect. Curr Biol 2007; 17(20):R877-8.
- 47. Motomura Y, Kitamura S, Oba K, Terasawa Y, Enomoto M, Katayose Y et al. Sleep debt elicits negative emotional reaction through diminished amygdala-anterior cingulate functional connectivity. PLoS One 2013; 8(2):e56578.
- 48. Goldstein AN, Walker MP. The role of sleep in emotional brain function. Annu Rev Clin Psychol 2014; 10:679–708.
- 49. Prather AA, Bogdan R, Hariri AR. Impact of sleep quality on amygdala reactivity, negative affect, and perceived stress. Psychosom Med 2013; 75(4):350–8.
- 50. Chuah LYM, Dolcos F, Chen AK, Zheng H, Parimal S, Chee MWL. Sleep deprivation and interference by emotional distracters. Sleep 2010; 33(10):1305–13.
- 51. Killgore WDS. Self-reported sleep correlates

with prefrontal-amygdala functional connectivity and emotional functioning. Sleep 2013; 36(11):1597–608.

- 52. Goldstein AN, Greer SM, Saletin JM, Harvey AG, Nitschke JB, Walker MP. Tired and apprehensive: anxiety amplifies the impact of sleep loss on aversive brain anticipation. J Neurosci 2013; 33(26):10607–15.
- Franzen PL, Buysse DJ, Dahl RE, Thompson W, Siegle GJ. Sleep deprivation alters pupillary reactivity to emotional stimuli in healthy young adults. Biol Psychol 2009; 80(3):300–5.
- 54. Goldstein-Piekarski AN, Greer SM, Saletin JM, Walker MP. Sleep Deprivation Impairs the Human Central and Peripheral Nervous System Discrimination of Social Threat. J Neurosci 2015; 35(28):10135–45.
- 55. Simon EB, Oren N, Sharon H, Kirschner A, Goldway N, Okon-Singer H et al. Losing Neutrality: The Neural Basis of Impaired Emotional Control without Sleep. J Neurosci 2015; 35(38):13194–205.
- 56. Alfarra R, Fins AI, Chayo I, Tartar JL. Changes in attention to an emotional task after sleep deprivation: neurophysiological and behavioral findings. Biol Psychol 2015; 104:1–7.
- 57. Mallick BN, Singh A. REM sleep loss increases brain excitability: role of noradrenaline and its mechanism of action. Sleep Med Rev 2011; 15(3):165–78.
- Siegel JM, Rogawski MA. A function for REM sleep: regulation of noradrenergic receptor sensitivity. Brain Res 1988; 472(3):213–33.