

The Role of Diet and Lifestyle in Preventing Oral Diseases: A Meta-Analysis

ARTICLE INFO

Article Type: Original Paper

Authors:

Hossein Malekzadeh¹ Amir Danial Pour Ahmadiyeh^{2*}

- Department of Oral and Maxillofacial Medicine, School of Dentistry, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.
- Department of Oral and Maxillofacial Surgery, School of Dentistry, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

* Corresponding author:

Dr. Amir Danial Pour Ahmadiyeh

E-mail: dr.ad.pourahmadie@gmail.com

ABSTRACT

Background: Oral health issues, such as dental caries, periodontal disease, and oral cancer, represent significant challenges to global health. Proper and balanced nutrition plays a vital role in this context, as it is fundamental to overall health, which includes the condition of the oral cavity and teeth. Nutrition pertains to the specific nutrients absorbed by the body, while diet encompasses all nutrients and non-nutrients consumed. The quality of nutrition directly influences the growth, development, and metabolic functions of the periodontium. A deficiency in essential nutrients can lead to changes in the primary factors associated with periodontal diseases, and dietary choices can significantly influence the progression of these ailments. This review article provides a meta-analysis of the existing literature to evaluate the effects of dietary and lifestyle factors on the prevention of oral health problems.

Method: A systematic review of studies was performed from databases such as PubMed, Scopus, Web of Science, and the Cochrane Library from 2000 onward to investigate the connections between specific dietary components, lifestyle habits, and oral health outcomes.

Results: The findings indicate notable correlations among elevated consumption of processed sugars, inadequate oral hygiene, and the use of tobacco and alcohol, all of which increase the likelihood of developing oral diseases. Conversely, a diet abundant in fruits, vegetables, and vital nutrients, along with consistent oral hygiene practices, is linked to a lower risk of these health issues.

Conclusion: This meta-analysis highlights the essential role that dietary and lifestyle modifications play in strategies aimed at preventing oral diseases.

Keywords:

Diet, Lifestyle, Nutrition, Oral Disease, Periodontal.

Copyright© 2020, TMU Press. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms.

INTRODUCTION

Oral diseases, although primarily preventable, represent a significant health challenge for nations, impacting individuals numerous throughout their lives and leading to pain, discomfort, disfigurement, and in severe cases, mortality (1). Recent estimates suggest that around 3.5 billion individuals are affected by oral health problems. The Global Burden of Disease 2021 report highlights untreated dental caries in permanent teeth as the most prevalent health issue (2). The management and prevention of oral health issues can lead to considerable expenses and are often excluded from the benefit packages of national universal health coverage (UHC) (3). In many low- and middle-income countries, there is a lack of adequate services to effectively prevent and manage oral health conditions (4). The causes of oral diseases are linked to several modifiable risk factors that are prevalent in numerous noncommunicable diseases (NCDs), such as excessive sugar intake, the use of tobacco and alcohol, and inadequate hygiene practices, as well as the associated social and commercial determinants (5). The lack of preventive measures in oral health significantly heightens the risk of tooth loss, which in turn diminishes an individual's capacity to chew and adversely affects their nutritional consumption. Furthermore, there are cosmetic consequences, as individuals with dental issues frequently refrain from smiling and may withdraw from social

interactions (6). The dental challenges faced by economically disadvantaged populations not only compromise their personal health but also place a financial burden on the overall healthcare system, as oral diseases can result in more severe health issues, leading individuals to pursue formal medical treatment at different stages of their lives (7). Certain lifestyle elements can increase the likelihood of developing periodontal disease, with inadequate oral hygiene serving as a significant factor; the absence of bacteria is essential for the development of periodontal disease, underscoring its importance in the condition's emergence (8). Efficient management of plaque, which involves the effective disruption of biofilm, is recognized as the most critical preventive strategy against periodontal diseases (9). Effective plaque control is largely determined by personal habits, making it an important aspect of one's lifestyle. Numerous studies carried out over the years have highlighted the significant impact of tobacco smoking on the onset of periodontitis through various mechanisms (10). These mechanisms include alterations in the oral microbiome that promote the growth of more harmful bacteria, reduced blood flow to the gums, impaired neutrophil function, elevated levels proinflammatory agents, and an increase in immunogenic T-cells (11). Furthermore, smoking adversely affects the recovery mechanisms of the impacted gingival tissues (12). Additionally, studies have indicated that the intake of alcohol correlates with a significant increase in the severity of periodontal disease, establishing it as a distinct modifiable risk factor for periodontitis (13-15). The underlying mechanisms suggest that high levels of alcohol consumption negatively impact the oral microbiome and interfere with the activities of neutrophils, macrophages, and Tcells. This disruption subsequently modifies immune responses and impedes the process of bone regeneration (16).

There exists a distinct psychobiological connection between chronic stress and associated health complications, particularly periodontal diseases. Chronic stress hampers tissue healing elevating the levels of cytokines, interleukins—especially IL-1β, IL-6, and IL-8 and TNF-α, which exacerbates periodontal damage (17).Furthermore. stress exerts considerable strain on multiple components of the immune system, including mitogen stimulation and the production of antibodies and cytokines, leading to an unbalanced immune response that facilitates the onset of periodontitis (18). Additionally, diets high in sugar or saturated fats, along with those deficient in polyols, fiber, and polyunsaturated fats, have been associated with a greater vulnerability to periodontal disease (19). consistently Research has demonstrated relationship between insufficient dietary calcium and the incidence of periodontal disease, with a notably low serum calcium-magnesium ratio being strongly associated with greater attachment and the advancement of periodontal conditions (21-22). As a result, dairy products, which are rich in calcium, phosphate, and a variety of proteins, are expected to positively influence periodontal health (22). In contrast, insufficient calcium consumption is linked to a heightened risk of tooth loss, loss of attachment, and increased severity of periodontal disease the current meta-analysis Therefore, highlights the crucial importance of dietary and lifestyle modifications in strategies designed to prevent oral diseases.

METHODS

Search Strategy

The comprehensive literature search in databases, including PubMed, Scopus, Web of Science, and the Cochrane Library, were utilized focusing on studies published from 2000 onward by employing search terms and keywords such as "dental caries," "periodontal disease," "oral cancer," "diet," "nutrition," "oral hygiene," "smoking," "alcohol," "lifestyle," "Systemic disease," "Stress," and "exercise" among others.

Inclusion and Exclusion Criteria

Numerous research methodologies, including randomized controlled trials, cohort studies, case-control studies, and observational studies, have been employed to explore the association between dietary elements, lifestyle choices, and oral health outcomes. It is essential that these studies include those that yield quantitative data relevant to the findings. Additionally, research involving non-human subjects, studies marked by insufficient data, case studies, literature reviews, and investigations lacking a direct link to dietary or lifestyle factors should be excluded from consideration.

Data Extraction

Two researchers conducted the data extraction process independently. The gathered data encompassed various elements, including the author's name, the country of origin, the total number of participants, the number of individuals at risk of malnutrition or already malnourished, the count of well-nourished participants, the type of study undertaken, the duration of follow-up, demographic information such as age and sex, oral health outcomes, the nutritional assessment scale employed, FTU, occurrences of oral disease, the mean DMFT, and classifications established by MNA/SGA.

Quality Assessment

The researchers, PSA and FWMGM, who were unaware of the study conditions, conducted independent evaluations to detect any possible biases present in the studies. They employed the Agency for Healthcare Research and Quality (AHRQ) scale, specifically designed for cross-sectional studies, alongside the Newcastle-Ottawa quality assessment scale, which is utilized for assessing case-control and cohort studies.

Statistical Analysis Method

The process of conducting a meta-analysis commenced with the extraction of statistical data from each relevant article, emphasizing the comparisons made among various groups. This data included the sample sizes for each group, the methods under comparison, and the means and standard deviations of the P-values, along with any additional pertinent statistical information provided by the authors. Following this, the homogeneity of the comparisons was evaluated through the chi-squared test. In instances where the studies exhibited homogeneity, fixed effect

models were utilized. The effect size for each comparison was determined according to the nature of the collected data; for example, when the response variable was minimal, the standardized difference in means was calculated. Ultimately, the cumulative effect was assessed based on the P-value, which informed the choice of the most appropriate method for each category of comparisons. If the comparisons within each category were homogeneous, the cumulative effect was computed using random effect models, supported by Comprehensive Meta-analysis software version 2.0.

RESULTS


Search Results

In this research, a comprehensive search was conducted across the PubMed, Scopus, Web of Science, and the Cochrane Library databases utilizing the specified keywords. Following the removal of duplicate entries, a total of 74 articles were identified according to the modifiable factors showed in Table 1. An initial review of the abstracts allowed for the elimination of unrelated studies, resulting in the preparation and examination of the full texts of 65 articles. No additional articles were discovered during the manual search of the references within these studies. From the 65 articles reviewed, those deemed relevant were chosen for a more in-depth analysis. This evaluation focused on two key aspects: the scientific principles underpinning the studies and the precision of their methodologies. Consequently, 25 articles were excluded, leaving 40 articles that were deemed suitable for statistical analysis. The PRISMA flowchart demonstrate the selection process (Table 2).

Table 1: Considered factors in the systematic review.

| Dietary Factors | | | | | | | |
|---|--|--|--|--|--|--|--|
| High sugar consumption and dental caries risk. | | | | | | | |
| Dietary fiber and dental caries risk | | | | | | | |
| Fermentable carbohydrates, acidity, and dental caries | | | | | | | |
| Fruit and vegetable consumption and periodontal health | | | | | | | |
| Calcium, Vitamin D, and periodontal health | | | | | | | |
| Antioxidants and oral cancer prevention | | | | | | | |
| Lifestyle Factors | | | | | | | |
| Oral hygiene practices (brushing frequency, flossing, etc.) and dental caries | | | | | | | |
| Oral hygiene practices and periodontal disease | | | | | | | |
| Smoking and periodontal disease, oral cancer | | | | | | | |
| Alcohol consumption and oral cancer | | | | | | | |
| Calcium, Vitamin D, and periodontal health | | | | | | | |
| Obesity and periodontal disease, if applicable | | | | | | | |
| Sstemic disease and stress | | | | | | | |

Table 2: Prisma chart of studies included in the systematic review.



Meta-Analysis of Studies

A meta-analysis was conducted once more to evaluate the variable of risk factors, as presented in Table 3. The findings from the assessment of risk factors revealed a notable distinction between groups, indicating that the life style and diet not only exhibited reduced oral disease but also experienced an increase in oral health (Table 4).

Table 3: Articles in which the mentioned factors in oral diseases were included in the meta-analysis.

| Nutrients | Author | Author Ref Results | | Lifestyle | Author | Ref | Results |
|-----------------|----------------------------|--------------------|--|--|---------------------------|-----|---|
| | Dodington (2015) | 24 | Improve outcomes of periodontal therapy | Weekly alcohol consumption | Baumeister (2021) | 44 | A risk factor for the development of periodontitis |
| | Neiva (2005) | 25 | Accelerate periodontal wound healing | Long-term alcohol intake | Pinto (2024) | 45 | Damaged periapical bone microstructure in the presence of apical periodontitis |
| Vitamins | Abou Sulaiman (2010) | 26 | Improvement of chronic periodontitis. | Excessive alcohol consumption | Moura (2024) | 46 | Increased osteoclast activity |
| | Aral (2015) | 27 | Alleviate gingival inflammation and alveolar bone | Appropriate physical exercise | Flynn (2007) | 47 | Reduce inflammation and osteoclast formation near the alveolar bone |
| Antioxidants | Grover (2016) | 28 | Reducing inflammation and periodontal destruction | Regular exercise | Shimazaki (2010) | 48 | Preventing periodontitis |
| | Kaur (2016) | 29 | Accelerate Periodontal regeneration | Moderate obstructive sleep apnea | Khodadadi (2022) | 49 | Increase risk of periodontitis |
| Magnesium | P Meisel (2005) | 30 | Improve periodontal health | Sleep duration of 7–9 h | Zhou (2021) | 50 | Lower oral health care needs |
| Probiotics | Morales (2017) | 31 | Supportive periodontal therapy. | High blood pressure | Zhan (2016) | 51 | Increase risk of periodontitis |
| Fungal extracts | Spratt | | Anticaries activity | Coronary heart diseases | Humphrey (2008) | 52 | Increase risk of periodontitis |
| Lycopene | Chandra (2007) | 22 | | Higher BMI | Sede (2014) | 53 | High caries incidence |
| Calcium | Mieko Nishida (2000) | 34 | Decreases tooth loss | Obesity | Abu- Shawish (2022) | 54 | Increase risk of periodontitis |

| Nutrients | Author | Ref | Results | Lifestyle | Author | Ref | Results | |
|-----------------------------|---------------------|-----|--|--|---------------------|-----|---|--|
| Melatonin | Arabacı (2015) | 35 | Reduces bone resorption caused by induced periodontitis | Obesity | Chen (2021) | 55 | Increase risk of chronic periodontitis | |
| Dietary acids | Carvalho (2020) | 36 | Association between dental erosion and the acidic foods | Former smoking/ current smoking | Khan (2016) | 56 | Periodontitis prevalence was highest in current smokers | |
| Starch-rich staple foods | Jangda (2024) | 37 | Low levels of caries | Uncontrolled diabetes | Braga (2011) | 57 | Accelerated orthodontic tooth movement | |
| Sucrose | Yoshihara (2021) | 38 | Rise in root surface caries. | Poor oral hygiene consumption | Najeeb (2017) | 58 | Affect periodontal tissues | |
| Fatty acids | Iwasaki (2011) | 39 | Incidence of periodontal disease events | Fitness | Moreno (2008) | 59 | Better oral health | |
| β-carotene | Iwasaki (2013) | 40 | Inverse correlation with the progression of periodontal disease. | Watching TV or working on a computer | Zeng (2014) | 60 | Increases of teeth cavities | |
| Milk | Adegboye (2012) | 41 | Reduced risk of developing periodontitis. | Regular tooth brushing | Hashmat (2023) | 61 | Decreases of caries rates | |
| Fruit juice | Liska (2019) | 42 | The erosion of dental enamel in adults. | work-related stress | Tikhonova (2018) | 62 | Increases risk of periodontal disease | |
| Polyunsaturated fatty acids | Iwasaki (2016) | 43 | Increased risk of hyposalivation | work stress | Sato (2020) | 63 | Increases tooth loss | |

 Table 4: Results of meta-analysis of studies.

| | Effect size and 95% confidence interval | | | | | Heterogeneity | | | Test of null (2-tail) | Tau-squared | | | |
|--------------|---|-------------------|-------------------|----------------|----------------|---------------|---------|----------|--------------------------|-----------------|-------------|----------------|-------|
| Model | Numbe r of Studies | Point estimate | Standard error | Upper bound | Lower bound | P-value | Q-value | Df (Q) | l-squared | <i>P</i> -value | Tau squared | Standard error | tau |
| cons tant | 10 | 0.031 | 0.011 | 0.044 | 0.002 | 0.213 | 2.98 | 11 0.000 | 0.167 | 0.000 | 0.0012 | 0.000 | |
| Ran dom | 6 | 0.028 | 0.015 | 0.053 | 0.001 | 0.119 | 2.50 | | 0.500 | 0.181 | 0.000 | 0.0012 | 0.000 |

Table 5. The quality of studies is evaluated in accordance with the Cochrane guidelines.

| Study | Random sequence generation | Allocation | Blinding | Incomplete data | Selective data | Free of bias |
|---------------|----------------------------|------------------|-------------------|------------------|------------------|-------------------|
| Dodington | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Neiva | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias |
| Abou Sulaiman | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias |
| Aral | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias |
| Grover | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| Kaur | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| P Meisel | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| Morales | Low risk of bias | Unrevealed | High risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| Spratt | Low risk of bias | Unrevealed | High risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| Chandra | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | High risk of bias |
| Mieko Nishida | Unrevealed | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | High risk of bias |
| Arabacı | Low risk of bias | Low risk of bias | High risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Carvalho | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias |
| Jangda | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Yoshihara | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Iwasaki | Unrevealed | Unrevealed | High risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |

| Study | Random sequence generation | Allocation | Blinding | Incomplete data | Selective data | Free of bias |
|------------|----------------------------|-------------------|-------------------|-------------------|------------------|------------------|
| Iwasaki | Unrevealed | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Adegboye | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Liska | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Iwasaki | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Baumeister | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Pinto | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias |
| Moura | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Flynn | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Shimazaki | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Khodadadi | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Zhou | Unrevealed | High risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| Zhan | Unrevealed | Low risk of bias | Low risk of bias | High risk of bias | Low risk of bias | Unrevealed |
| Humphrey | Unrevealed | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias |
| Sede | Unrevealed | Low risk of bias | Low risk of bias | High risk of bias | Low risk of bias | Low risk of bias |
| AbuShawish | Low risk of bias | Unrevealed | High risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| Chen | Unrevealed | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Unrevealed |
| Khan | Unrevealed | High risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed |
| Braga | Unrevealed | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Najeeb | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias |
| Moreno | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias |
| Zeng | Low risk of bias | Unrevealed | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias |
| Hashmat | Unrevealed | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed | Low risk of bias |
| Tikhonova | Low risk of bias | Unrevealed | Unrevealed | High risk of bias | Low risk of bias | Low risk of bias |
| Sato | Low risk of bias | Low risk of bias | Low risk of bias | Low risk of bias | Unrevealed | Unrevealed |

DISCUSSION

The present review article highlights the vital importance of diet in maintaining oral health, affecting both the onset and prevention of numerous dental issues. Certain dietary elements have been recognized as key factors in the progression of dental caries, periodontal disease, and oral cancers. Dental caries, often referred to as tooth decay, predominantly results from the erosion of tooth enamel caused by acids produced by bacteria in the mouth (64). The regular intake of refined sugars, such as sucrose and fructose, with fermentable carbohydrates like starches, notably heightens the likelihood of developing dental caries (65). These compounds act as a food source for oral bacteria, including Streptococcus mutans, which convert them into acids that lead to the demineralization of tooth enamel (66). Moreover, frequent snacking and the consumption of sugary drinks can intensify this problem. Additionally, the intake of acidic foods and beverages, including citrus fruits, soft drinks, and energy drinks, can lead to enamel erosion, which further threatens the structural integrity of the teeth (67).

Dietary fiber, especially soluble fiber, is essential for enhancing saliva production, which acts as a natural defense against acids and assists in the removal of plaque. Therefore, it is recommended to maintain a diet rich in whole grains, fruits, and vegetables to support optimal oral health (68). Additionally, fluoride, found in drinking water, toothpaste, and numerous other products, is vital for the remineralization of tooth enamel, thereby bolstering its resistance to acid erosion (69). Although certain foods may have trace amounts of fluoride, they are typically not the main contributors to the fluoride needed for maintaining oral health. Periodontal disease, which impacts the gums and the structures that support the teeth, mainly develops due to the buildup of bacterial plaque and the resulting inflammation. Consuming a diet abundant in antioxidants and anti-inflammatory substances can promote gum health and mitigate the inflammation linked to periodontal disease (70). Nutrient-dense foods, including vegetables—particularly leafy greens—omega-3 fatty acid-rich fish, and nuts, play a significant role in enhancing a robust immune response. Furthermore, sufficient intake of calcium and vitamin D is crucial for preserving bone density, which encompasses the alveolar bone that underpins the teeth, thereby promoting the overall health of both the bone and the soft tissues within the oral cavity (71).

A nutritional approach that prioritizes the

intake of whole, unrefined foods over highly processed options has demonstrated positive effects on systemic inflammation and has been linked to enhanced overall oral health. Dietary selections play a crucial role in the incidence of oral cancers, encompassing malignancies of the lip, tongue, mouth, and oropharynx (72). A significant consumption of processed foods and red meat is frequently associated with an increased risk of several types of cancer, including those that impact the oral cavity. In contrast, diets abundant in fruits and vegetables, which are rich in antioxidants and beneficial phytonutrients, are linked to a lower likelihood of developing oral cancers. Additionally, deficiencies in specific micronutrients, such as vitamins A, C, and E, as well as folate, have been correlated with an elevated susceptibility to oral cancer (73).

Lifestyle choices play a crucial role in determining oral health outcomes, impacting them both on their own and in conjunction with dietary practices. Adhering to appropriate oral hygiene routines is vital for the prevention of oral diseases. Regularly brushing teeth, a minimum of twice daily with fluoridated toothpaste is necessary to efficiently eliminate plaque and bacteria from the surfaces of the teeth (74). The proper technique for brushing and allowing sufficient time for this activity are essential components of an effective oral hygiene routine. Furthermore, the use of dental floss or similar interdental cleaning tools is vital for removing plaque and food debris from spaces between teeth toothbrushes are unable to Incorporating antiseptic mouthwashes can also enhance the benefits of regular brushing and flossing, as they assist in controlling bacterial populations and reducing the risk of gum disease Regular dental examinations (75).professional cleanings conducted by a dentist or dental hygienist are essential for the removal of tartar and the early identification of potential health concerns (76). The use of tobacco in any form significantly elevates the risk of developing oral cancers and periodontal disease. Both smoking and smokeless tobacco impair blood circulation and promote inflammation, which increases the likelihood of infections and impedes tissue healing, adversely affecting both soft tissues and bone structures within the oral cavity. Furthermore, excessive alcohol intake heightens

the risk of oral cancers, especially when combined with smoking, and disrupts the balance of the oral environment. Heavy alcohol consumption can damage the oral mucosa, rendering the oral cavity more susceptible to cellular damage and various diseases (77).

Chronic stress and insufficient sleep are associated with diminished immune function. thereby increasing the risk of inflammatory conditions, including periodontal disease. The presence of psychological stress leads to elevated levels of cortisol and inflammatory cytokines, both of which contribute to a heightened susceptibility to chronic inflammation and the development of periodontal disease Insufficient sleep also leads to increased cortisol levels and weakened immune function, which adversely affects overall health. Engaging in regular physical activity promotes overall wellbeing and can enhance oral health by improving reducing blood circulation. systemic inflammation, and aiding in stress management (79). Consistent exercise is instrumental in modulating inflammatory markers, mitigating stress, and can directly enhance oral health by reducing the overall burden of disease.

The interplay between diet, lifestyle, and oral health is intricate and involves numerous components. Various dietary and lifestyle factors frequently interact, affecting oral health results through different mechanisms. For instance, individuals who frequently ingest foods and beverages high in sugar while disregarding adequate dental hygiene face a significantly elevated risk of developing dental caries. Likewise, individuals who smoke or engage in excessive alcohol consumption are at a greater risk of oral cancer when compared to nonsmokers and those who drink in moderation (80). The distinct genetic makeup, socioeconomic status, and personal lifestyle choices of individuals significantly influence their oral health experiences (81). Public health initiatives play a vital role in raising awareness within the regarding the significance community maintaining a balanced diet, practicing regular oral hygiene, and steering clear of detrimental substances. It is imperative that these initiatives are customized to cater to various demographic segments and employ effective communication methods to inspire positive behavioral changes. Moreover, policy measures such as imposing

taxes on sugary beverages, regulating the marketing of tobacco products, and incorporating oral health education into school curricula can greatly improve public health and promote healthier lifestyle choices among community members. Healthcare professionals should focus on developing personalized prevention strategies by evaluating individual risk factors, dietary habits, and lifestyle decisions while providing tailored advice. Prioritizing early intervention in children is essential for instilling effective oral hygiene practices. Additionally, ensuring that affordable dental care, including preventive services like cleanings, is readily available is crucial for enhancing oral health and decreasing the prevalence of oral diseases within the community.

CONCLUSION

The data unequivocally indicates that nutrition and lifestyle choices are essential in preventing oral health issues. Individuals can greatly diminish their likelihood of experiencing oral diseases and sustain excellent oral health over their lifetimes by embracing a nutritious diet, maintaining proper oral hygiene, steering clear of tobacco and excessive alcohol use, managing stress effectively, and participating in regular physical exercise. Ongoing research, education, and the development of policies are crucial for enhancing oral health results and alleviating the global impact of oral diseases. This proactive strategy necessitates cooperation among various healthcare sectors and community partners.

ACKNOWLEDGMENTS

This study was self-supported.

CONFLICT OF INTEREST

There is no conflict of interest.

REFERENCES

- 1. World Health Organization. Global oral health status report: towards universal health coverage for oral health by 2030. Executive summary. World Health Organization; 2022 Nov 18.
- 2. Wen PY, Chen MX, Zhong YJ, Dong QQ, Wong HM. Global burden and inequality of dental caries, 1990 to 2019. Journal of dental research. 2022 Apr;101(4):392-9.
- 3. Jadidfard MP, Tahani B. Painless cost

- control as a central strategy for universal oral health coverage: A critical review with policy guide. International Journal of Dental Hygiene. 2024 May 19.
- 4. Watt RG, Daly B, Allison P, Macpherson LM, Venturelli R, Listl S, Weyant RJ, Mathur MR, Guarnizo-Herreño CC, Celeste RK, Peres MA. Ending the neglect of global oral health: time for radical action. The Lancet. 2019 Jul 20;394(10194):261-72.
- 5. Wolf TG, Cagetti MG, Fisher JM, Seeberger GK, Campus G. Non-communicable diseases and oral health: an overview. Frontiers in oral health. 2021 Sep 3;2:725460.
- 6. Antoniadou M, Varzakas T. Breaking the vicious circle of diet, malnutrition and oral health for the independent elderly. Critical Reviews in Food Science and Nutrition. 2021 Oct 28;61(19):3233-55.
- 7. Gil-Montoya JA, Ferreira de Mello AL, Barrios R, Gonzalez-Moles MA, Bravo M. Oral health in the elderly patient and its impact on general well-being: a nonsystematic review. Clinical interventions in aging. 2015 Feb 11:461-7.
- 8. Di Stefano M, Polizzi A, Santonocito S, Romano A, Lombardi T, Isola G. Impact of oral microbiome in periodontal health and periodontitis: a critical review on prevention and treatment. International journal of molecular sciences. 2022 May 5;23(9):5142.
- 9. Lazar V, Ditu LM, Curutiu C, Gheorghe I, Holban A, Popa M, Chifiriuc C. Impact of dental plaque biofilms in periodontal disease: Management and future therapy. Periodontitis: A Useful Reference; Arjunan, P., Ed.; InTech Open: London, UK. 2017 Nov 15:11-42.
- 10. da Silva RV, Rangel TP, Corrêa MG, de Freitas Monteiro M, Casati MZ, Ruiz KG, Sallum EA, Casarin RC, Sallum AW. Smoking negatively impacts the clinical, microbiological, and immunological treatment response of young adults with Grade C periodontitis. Journal of Periodontal Research. 2022 Dec;57(6):1116-26.
- Vitkov L, Muñoz LE, Schoen J, Knopf J, Schauer C, Minnich B, Herrmann M, Hannig M. Neutrophils orchestrate the periodontal

- pocket. Frontiers in Immunology. 2021 Nov 24;12:788766.
- 12. Apatzidou DA. The role of cigarette smoking in periodontal disease and treatment outcomes of dental implant therapy. Periodontology 2000. 2022 Oct;90(1):45-61.
- 13. Alves-Costa S, Leite FR, Ladeira LL, Lima-Soares F, de Andrade Paes AM, de Souza BF, Nascimento GG, Ribeiro CC. Behavioral and metabolic risk factors associated with periodontitis in Brazil, 1990–2019: a multidimensional analysis for the Global Burden of Disease Study 2019. Clinical Oral Investigations. 2023 Dec;27(12):7909-17.
- 14. Plachokova AS, Gjaltema J, Hagens ER, Hashemi Z, Knüppe TB, Kootstra TJ, Visser A, Bloem BR. Periodontitis: A Plausible Modifiable Risk Factor for Neurodegenerative Diseases? A Comprehensive Review. International journal of molecular sciences. 2024 Apr 19;25(8):4504.
- 15. Wang J, Lv J, Wang W, Jiang X. Alcohol consumption and risk of periodontitis: A meta-analysis. Journal of clinical periodontology. 2016 Jul;43(7):572-83.
- 16. Knight ET, Liu J, Seymour GJ, Faggion Jr CM, Cullinan MP. Risk factors that may modify the innate and adaptive immune responses in periodontal diseases. Periodontology 2000. 2016 Jun;71(1):22-51.
- 17. Gunepin M, Derache F, Trousselard M, Salsou B, Risso JJ. Impact of chronic stress on periodontal health. Journal of Oral Medicine and Oral Surgery. 2018 Jan 1;24(1):44-50.
- 18. Boia S, Stratul ŞI, Boariu M, Ursoniu S, Goţia SL, Boia ER, Borza C. Evaluation of antioxidant capacity and clinical assessment of patients with chronic periodontitis treated with non-surgical periodontal therapy and adjunctive systemic antibiotherapy. Romanian J Morphol Embryol. 2018 Jan 1;59(4):1107-13.
- 19. Martinon P, Fraticelli L, Giboreau A, Dussart C, Bourgeois D, Carrouel F. Nutrition as a key modifiable factor for periodontitis and main chronic diseases. Journal of clinical medicine. 2021 Jan 7;10(2):197.

- 20. Tanaka K, Miyake Y, Okubo H, Hanioka T, Sasaki S, Miyatake N, Arakawa M. Calcium intake is associated with decreased prevalence of periodontal disease in young Japanese women. Nutrition journal. 2014 Dec;13:1-6.
- 21. Varela-López A, Giampieri F, Bullón P, Battino M, Quiles JL. A systematic review on the implication of minerals in the onset, severity and treatment of periodontal disease. Molecules. 2016 Sep 7;21(9):1183.
- 22. Najeeb S, Zafar MS, Khurshid Z, Zohaib S, Almas K. The role of nutrition in periodontal health: an update. Nutrients. 2016 Aug 30;8(9):530.
- 23. Meisel P, Pink C, Nauck M, Jablonowski L, Voelzke H, Kocher T. Magnesium/calcium ratio in serum predicts periodontitis and tooth loss in a 5-year follow-up. JDR Clinical & Translational Research. 2016 Oct;1(3):266-74.
- 24. Dodington DW, Fritz PC, Sullivan PJ, Ward WE. Higher intakes of fruits and vegetables, β-carotene, vitamin C, α-tocopherol, EPA, and DHA are positively associated with periodontal healing after nonsurgical periodontal therapy in nonsmokers but not in smokers. The Journal of nutrition. 2015 Nov 1;145(11):2512-9.
- 25. Neiva RF, Al-Shammari K, Nociti Jr FH, Soehren S, Wang HL. Effects of vitamin-B complex supplementation on periodontal wound healing. Journal of periodontology. 2005 Jul;76(7):1084-91.
- 26. Abou Sulaiman AE, Shehadeh RM. Assessment of total antioxidant capacity and the use of vitamin C in the treatment of nonsmokers with chronic periodontitis. Journal of periodontology. 2010 Nov;81(11):1547-54.
- 27. Aral K, Alkan BA, Saraymen R, Yay A, Şen A, Önder GÖ. Therapeutic effects of systemic vitamin k2 and vitamin d3 on gingival inflammation and alveolar bone in rats with experimentally induced periodontitis. Journal of periodontology. 2015 May;86(5):666-73..
- 28. Grover V, Malhotra R, Kapoor A, Bither R, Sachdeva S. Correlation of alkaline

- phosphatase activity to clinical parameters of inflammation in smokers suffering from chronic periodontitis. Journal of Indian Society of Periodontology. 2016 May 1;20(3):254-9...
- 29. Muwal P, Kaur N, Kaur G. DUAL ROLE OF AUTOPHAGY IN PERIODONTAL DISEASE. infection.;8(9):10..
- 30. Meisel P, Schwahn C, Luedemann J, John U, Kroemer HK, Kocher T. Magnesium deficiency is associated with periodontal disease. Journal of dental research. 2005 Oct;84(10):937-41..
- 31. Morales AB, Bedoya J, Gamonal J. Probiotics and Periodontal diseases. Insights into various aspects of oral health. 2017 Sep 20:73-96..
- 32. Spratt DA, Daglia M, Papetti A, Stauder M, O' Donnell D, Ciric L, Tymon A, Repetto B, Signoretto C, Houri-Haddad Y, Feldman M. Evaluation of plant and fungal extracts for their potential antigingivitis and anticaries activity. BioMed Research International. 2012;2012(1):510198...
- 33. Chandra RV, Prabhuji ML, Roopa DA, Ravirajan S, Kishore HC. Efficacy of lycopene in the treatment of gingivitis: a randomised, placebo-controlled clinical trial. Oral health & preventive dentistry. 2007 Sep 1;5(4)..
- 34. Nishida M, Grossi SG, Dunford RG, Ho AW, Trevisan M, Genco RJ. Calcium and the risk for periodontal disease. Journal of periodontology. 2000 Jul;71(7):1057-66..
- 35. Arabacı T, Kermen E, Özkanlar S, Köse O, Kara A, Kızıldağ A, Duman ŞB, Ibişoğlu E. Therapeutic effects of melatonin on alveolar bone resorption after experimental periodontitis in rats: a biochemical and immunohistochemical study. Journal of periodontology. 2015 Jul;86(7):874-81.
- 36. Carvalho TS, Lussi A. Acidic beverages and foods associated with dental erosion and erosive tooth wear. The impact of nutrition and diet on oral health. 2020;28:91-8..
- 37. Jangda FH, Suominen AL, Lundqvist A, Männistö S, Golkari A, Bernabé E. Starch intake and caries increment: A longitudinal

- study in Finnish adults. Public Health Nutrition. 2024 Jan;27(1):e263..
- 38. Yoshihara A, Suwama K, Miyamoto A, Watanabe R, Ogawa H. Diet and root surface caries in a cohort of older Japanese. Community Dentistry and Oral Epidemiology. 2021 Jun;49(3):301-8..
- 39. Iwasaki M, Taylor GW, Moynihan P, Yoshihara A, Muramatsu K, Watanabe R, Miyazaki H. Dietary ratio of n-6 to n-3 polyunsaturated fatty acids and periodontal disease in community-based older Japanese: a 3-year follow-up study. Prostaglandins, leukotrienes and essential fatty acids. 2011 Aug 1;85(2):107-12..
- 40. Iwasaki M, Moynihan P, Manz MC, Taylor GW, Yoshihara A, Muramatsu K, Watanabe R, Miyazaki H. Dietary antioxidants and periodontal disease in community-based older Japanese: a 2-year follow-up study. Public health nutrition. 2013 Feb;16(2):330-8...
- 41. Adegboye AR, Christensen LB, Holm-Pedersen P, Avlund K, Boucher BJ, Heitmann BL. Intake of dairy products in relation to periodontitis in older Danish adults. Nutrients. 2012 Sep 4;4(9):1219-29...
- 42. Liska D, Kelley M, Mah E. 100% fruit juice and dental health: A systematic review of the literature. Frontiers in public health. 2019 Jul 12;7:190..
- 43. Iwasaki M, Yoshihara A, Ito K, Sato M, Minagawa K, Muramatsu K, Watanabe R, Manz MC, Ansai T, Miyazaki H. Hyposalivation and dietary nutrient intake among community-based older J apanese. Geriatrics & gerontology international. 2016 Apr;16(4):500-7..
- 44. Baumeister SE, Freuer D, Nolde M, Kocher T, Baurecht H, Khazaei Y, Ehmke B, Holtfreter B. Testing the association between tobacco smoking, alcohol consumption, and risk of periodontitis: A Mendelian randomization study. Journal of clinical periodontology. 2021 Nov;48(11):1414-20...
- 45. de Sousa Pinto CP. *Periodontal treatment impact on halitosis and quality of life* (Doctoral dissertation, Universidade do Porto (Portugal))..

- 46. de Moura JD, de Almeida Rodrigues P, Dos Santos VR, Bittencourt LO, Matos-Sousa JM, Peinado BR, Perdigão JM, Rogez H, Collares FM, Lima RR. Açai (Euterpe oleracea Mart.) supplementation promotes histological and ultrastructural changes in rats' alveolar bone. Heliyon. 2024 May 30;10(10)...
- 47. Flynn MG, McFarlin BK, Markofski MM. State of the art reviews: inflammatory actions of exercise training. American journal of lifestyle medicine. 2007 May;1(3):220-35...
- 48. Shimazaki Y, Egami Y, Matsubara T, Koike G, Akifusa S, Jingu S, Yamashita Y. Relationship between obesity and physical periodontitis. fitness and Journal periodontology. 2010 Aug;81(8):1124-31.
- 49. Khodadadi N, Khodadadi M, Zamani M. Is periodontitis associated with obstructive sleep apnea? A systematic review and meta-Journal analysis. of Clinical and Experimental Dentistry. 2022 Apr;14(4):e359...
- 50. Zhou F, Liu Z, Guo Y, Xu H. Association of short sleep with risk of periodontal disease: meta-analysis and Mendelian randomization study. Journal of Clinical Periodontology. 2021 Aug;48(8):1076-84...
- 51. Zhan Y, Jiao J, Jing W, Feng X, Tai B, Hu D, Lin HC, Wang B, Wang C, Zheng S, Liu X. between Association periodontitis hypertension: cross-sectional survey from the Fourth National Oral Health Survey of China open. (2015-2016).**BMJ** 2023 Mar 1;13(3):e068724...
- 52. Humphrey LL, Fu R, Buckley DI, Freeman M, Helfand M. Periodontal disease and coronary heart disease incidence: systematic review and meta-analysis. Journal general internal medicine. 2008 Dec;23:2079-86..
- 53. Sede MA, Ehizele AO. Relationship between obesity and oral diseases. Nigerian journal of clinical practice. 2014;17(6):683-90..
- 54. Abu-Shawish G, Betsy J, Anil S. Is obesity a risk factor for periodontal disease in adults? A systematic review. International Journal of Environmental Research and Public Health.

- 2022 Oct 4;19(19):12684...
- 55. Chen TP, Yu HC, Lin TH, Wang YH, Chang YC. Association between obesity and chronic periodontitis: A nationwide population-based cohort study in Taiwan. Medicine. 2021 Oct 15;100(41):e27506...
- 56. Khan S, Khalid T, Awan KH. Chronic periodontitis and smoking Prevalence and dose-response relationship. Saudi medical journal. 2016 Aug;37(8):889..
- 57. Braga SM, de Albuquerque Taddei SR, Andrade Jr I, Queiroz-Junior CM, Garlet GP, Repeke CE, Teixeira MM, da Silva TA. Effect of diabetes on orthodontic tooth movement in a mouse model. European Journal of Oral Sciences. 2011 Feb;119(1):7-
- 58. Najeeb S, Siddiqui F, Qasim SB, Khurshid Z, Zohaib S. Zafar MS. Influence uncontrolled diabetes mellitus on periodontal tissues during orthodontic tooth movement: a systematic review of animal studies. Progress in Orthodontics. 2017 Dec;18:1-7...
- 59. Moreno-Sandoval A, Guirao JM. Morphosyntactic tagging of the Spanish C-ORAL-ROM corpus: methodology, tools and evaluation. InSpoken language corpus and linguistic informatics 2008 Jul 1 (pp. 199-218). John Benjamins Publishing Company..
- 60. Zeng X, Sheiham A, Sabbah W. The association between dental caries television viewing among Chinese adolescents in Guangxi, China. BMC oral health. 2014 Dec;14:1-6...
- 61. Hashmat, S., Hingorjo, M.R., Shahid, R.A., Ghani, R., Rashid, M.N., Butt, F. (2023). Relationship between lifestyle and oral health. Biol. Clin. Sci. Res. J., 2023: 522. doi:
 - https://doi.org/10.54112/bcsrj.v2023i1.522].
- 62. Tikhonova S, Booij L, D'Souza V, Crosara KT, Siqueira WL, Emami E. Investigating the association between stress, saliva and dental caries: a scoping review. BMC oral health. 2018 Dec;18:1-9...
- 63. Sato Y, Saijo Y, Yoshioka E. Work stress and oral conditions: a systematic review of observational studies. BMJ open. 2021 May

- 1;11(5):e046532...
- 64. Saidaiah P, Banu Z, Khan AA, Geetha A, Somraj B. A comprehensive review of omega-3 fatty acids: Sources, industrial applications, and health benefits. Annals of Phytomedicine. 2024;13(1):209-25.
- 65. Botelho J, Machado V, Proença L, Delgado AS, Mendes JJ. Vitamin D deficiency and oral health: a comprehensive review. Nutrients. 2020 May 19;12(5):1471.
- 66. Rodríguez-Molinero J, Migueláñez-Medrán BD, Puente-Gutiérrez C, Delgado-Somolinos E, Martin Carreras-Presas C, Fernández-Farhall J, López-Sánchez AF. Association between oral cancer and diet: An update. Nutrients. 2021 Apr 15;13(4):1299.
- 67. Fagbohun OF, Gillies CR, Murphy KP, Rupasinghe HV. Role of antioxidant vitamins and other micronutrients on regulations of specific genes and signaling pathways in the prevention and treatment of cancer. International Journal of Molecular Sciences. 2023 Mar 23;24(7):6092.
- 68. Ramakrishnan M, Shukri MM. Fluoride, Fluoridated Toothpaste Efficacy And Its Safety In Children-Review. International Journal of Pharmaceutical Research (09752366). 2018 Oct 1;10(4).
- 69. Ahuja A, Singh S. Impact of the current scenario and future perspectives for the management of oral diseases: Remarkable contribution of herbs in dentistry. Anti-Infective Agents. 2022 Oct 1;20(5):27-45.
- 70. Zhang Y, He J, He B, Huang R, Li M. Effect of tobacco on periodontal disease and oral cancer. Tobacco induced diseases. 2019;17.
- 71. Pelucchi C, Gallus S, Garavello W, Bosetti C, La Vecchia C. Cancer risk associated with alcohol and tobacco use: focus on upper aero-digestive tract and liver. Alcohol Research & Health. 2006;29(3):193.
- 72. Decker A, Askar H, Tattan M, Taichman R, Wang HL. The assessment of stress, depression, and inflammation as a collective risk factor for periodontal diseases: a systematic review. Clinical oral investigations. 2020 Jan;24:1-2.
- 73. Barranca-Enríquez A, Romo-González T.

- Your health is in your mouth: A comprehensive view to promote general wellness. Frontiers in Oral Health. 2022 Sep 14;3:971223.
- 74. Tran Q, Maddineni S, Arnaud EH, Divi V, Megwalu UC, Topf MC, Sunwoo JB. Oral cavity cancer in young, non-smoking, and non-drinking patients: A contemporary review. Critical reviews in oncology/hematology. 2023 Oct 1;190:104112.
- 75. Patrick DL, Lee RS, Nucci M, Grembowski D, Jolles CZ, Milgrom P. Reducing oral health disparities: a focus on social and cultural determinants. InBMC oral health 2006 Jun (Vol. 6, pp. 1-17). BioMed Central.
- 76. Abdelaziz M. Detection, Diagnosis, and Monitoring of Early Caries: The Future of Individualized Dental Care. Diagnostics. 2023 Dec 12;13(24):3649.
- 77. Reidy J, McHugh E, Stassen LF. A review of the relationship between alcohol and oral cancer. The surgeon. 2011 Oct 1;9(5):278-83.
- 78. Decker A, Askar H, Tattan M, Taichman R, Wang HL. The assessment of stress, depression, and inflammation as a collective risk factor for periodontal diseases: a systematic review. Clinical oral investigations. 2020 Jan;24:1-2.
- 79. Abe M, Abe H. Lifestyle medicine—An evidence based approach to nutrition, sleep, physical activity, and stress management on health and chronic illness. Personalized Medicine Universe. 2019 Jul 1;8:3-9.
- 80. Jaber MA, Porter SR, Scully C, Gilthorpe MS, Bedi R. The role of alcohol in non-smokers and tobacco in non-drinkers in the aetiology of oral epithelial dysplasia. International journal of cancer. 1998 Jul 29;77(3):333-6.
- 81. Heilmann A, Tsakos G, Watt RG. Oral health over the life course. A life course perspective on health trajectories and transitions. 2015:39-59.