



Anti-Inflammatory Reflex - Interaction of The Nervous System and The Immune System

ARTICLE INFO

Article Type

Narrative Review

Author

Roham Mazloom

Department of Physiology-Pharmacology-
Medical Physics, School of Medicine,
Alborz University of Medical Sciences,
Karaj, Iran.

*Corresponding author:

Roham Mazloom,

Department of Physiology-Pharmacology-
Medical Physics, Faculty of Medicine,
Alborz University of Medical Sciences,
Karaj, Iran.

rohammazloom@abzums.ac.ir

Fax: +98-2634287339

ABSTRACT

The coordination of body systems is important for overcoming various conditions. The nervous system, as a fast coordinator of the body, reflexively manages many functions. At the same time, the immune system is involved in endogenous and exogenous factors that disturb homeostasis at any time to protect the body. Research has shown negative feedback between the two systems. Inflammatory factors such as pro-inflammatory cytokines stimulate the vagus nerve, which increases afferent signals to the central nervous system. The central autonomic network increases efferent vagus nerve impulses. Amplification of efferent vagus nerve activity promotes the release of acetylcholine. Increased acetylcholine suppresses inflammation through its receptors on immune cells. The aforementioned feedback process, which is the two-way communication of the nervous and immune systems, is called the "anti-inflammatory reflex". In the present article, the role of each component and the therapeutic potential of using the anti-inflammatory reflex will be discussed. Moreover, heart rate variability as an index for measuring the state of the anti-inflammatory reflex is considered.

Keywords: anti-inflammatory reflex; inflammation; autonomic nervous system; vagus nerve; immune system

Copyright© 2020, TMU Press. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms

INTRODUCTION

In biology, various nervous reflexes unconsciously manage many body functions using afferent and efferent neurons [1]. The receptors of the afferent neurons are activated by stimuli. Activation of these receptors increases the impulses of afferent neurons in the central nervous system. The central nervous system processes input neural information and sends back responses through efferent neurons. Upon each reflex, a specific stimulus can affect receptors in the afferent neurons, which triggers circuits [2, 3]. Simultaneously, the body faces certain conditions that can affect all systems, including inflammatory processes.

Inflammation is the body's protective response to injury or infection and is accompanied by swelling, pain, redness, and warmth [4, 5]. Inflammatory diseases are diverse, and there is no

specific treatment for them under chronic conditions [6, 7]. During the process of inflammation, the immune cells secrete cytokines, which are divided into two types, pro-inflammatory and anti-inflammatory [8, 9]. The goal is to secrete cytokines that limit inflammation [10]. However, the intense or continuous activity of macrophages leads to excessive production of cytokines, causing them to escape local control and enter the blood circulation system, which widely activates the inflammatory component, systemic inflammatory response syndrome (SIRS) [11-14].

Previous studies have indicated an interaction between the nervous system and inflammation [15, 16]. Among the various parts of the nervous system, the vagus nerve plays the most important role in anti-inflammatory reflexes owing to its extensiveness. Borovikova et al. were the first to

formally describe the role of vagal stimulation in reducing the inflammatory response caused by endotoxins [17]. Subsequently, Kevin J. Tracey proposed a reflex that is activated by inflammation and controls it [15]. Various studies over the last two decades have confirmed the Tracey hypothesis. Inflammation activates the afferent vagus nerve, which triggers the central autonomic network (CAN). Evoked CAN increased efferent autonomic nerve activity, which elevated acetylcholine content. High acetylcholine content modulates inflammation through interactions with immune cells. This entire process is called the “cholinergic inflammatory reflex.” Many studies have shown that different inflammatory conditions activate the anti-inflammatory reflex [18-20]. To better define the anti-inflammatory reflex pattern, the available evidence for each stage will be reviewed separately.

Stimulation of the afferent vagus nerve under inflammation

The vagus nerve, which is widely distributed in visceral areas, consists of more afferent fibers than efferent fibers [21]. Previous investigations indicated that stimulation of the vagus nerve is the first step of the anti-inflammatory reflex [22, 23]. Toll-like receptors (TLRs) are critical for triggering the innate immune response to defend against infections [24]. The components of Gram-negative and Gram-positive bacteria interact with TLR4 and TLR2, respectively, in monocytes and macrophages, which release different chemokines and cytokines. These cytokines recruit and activate neutrophils to eliminate pathogens [25, 26]. Pathogen-associated molecular patterns (PAMPs) and pro-inflammatory cytokines such as TNF- α , IL-6, high mobility group box 1 (HMGB1), and IL-1 β , which are released under various inflammatory conditions, can stimulate the vagus nerve [27, 28]. It seems that pro-inflammatory cytokines interact with their receptors in the afferent vagus nerve [29]. Moreover, direct stimulation of the vagus nerve triggers an anti-inflammatory reflex [30]. Altogether, the elevation of afferent vagus nerve activity alters CAN.

CAN processing of increased vagus nerve activity under inflammation

The first part of the CAN that receives afferent messages from the vagus nerve is the nucleus tractus solitaries (NTS) [31]. As a hub, the NTS sends the desired messages received from the vagus nerve to the rest of the CAN and other parts of the central nervous system [32]. Different key components are observed in CAN, including the insular cortex, amygdala, hypothalamus, periaqueductal gray matter, parabrachial complex, NTS, and ventrolateral medulla [33]. Finally, after processing, the efferent output of the vagus nerve originates from the dorsal motor nucleus of the vague (DMNV) [34]. Moreover, the direct stimulation of some neuronal pathways in the CNS can increase afferent vagus nerve signals. For example, activation of the cholinergic brain network through muscarinic agonists augments efferent vagus nerve signals [35]. Furthermore, the areas of CAN are known components of higher brain function [33, 36]. Higher brain functions such as cognition and emotion interact with CAN which alters vagus nerve activity [37]. Therefore, CAN as a processor combines different signals and finally changes the activity level of the vagus nerve.

Increasing the efferent vagus nerve activity

Extensive research has been conducted on the anti-inflammatory effects of increasing the excitatory responses of the vagus nerve. Stimulation of the vagus nerve can directly elevate the amount of acetylcholine secretion [38, 39]. Increasing the secretion of vagal acetylcholine affects various receptors in the areas innervated by it, such as the heart, intestine, stomach, and liver [35, 40]. Research has shown that a type of nicotinic acetylcholine receptor can induce anti-inflammatory processes. Acetylcholine magnification affects alpha7-nicotinic acetylcholine receptors (α_7 nAChR) in immune cells, including macrophages, and prevents the release of more pro-inflammatory cytokines, resulting in inflammation control [41-43]. α_7 nAChR is present in many tissues of the body and is not limited to the nervous system, but with its effect, it can help limit inflammation through immune cells [44, 45].

Moreover, a branch of the autonomic nerve is sent to the celiac ganglion, the postganglionic splanchnic extension of which releases norepinephrine in the spleen. Norepinephrine release in the spleen interacts with beta2-adrenergic receptors (β 2AR) in CD4 + T-lymphocyte cells [46]. This type of cell contains choline acetyltransferase (ChAT), which reacts with norepinephrine by releasing acetylcholine [47, 48]. Acetylcholine produced by splenic T-lymphocyte cells can act on α 7nAChR in the spleen and other tissues, thereby reducing inflammation [49].

Hypothalamus-Pituitary-Adrenal axis on the management of inflammation

In addition to the above-mentioned processes, the hypothalamus-pituitary-adrenal (HPA) axis can reduce inflammation [50]. This pathway aligns with the anti-inflammatory reflex. Therefore, stimulation from the NTS to the hypothalamus increases the production and release of corticotropin-releasing hormone (CRH). The effect of CRH on the anterior pituitary gland is to increase the release of adrenocorticotropic hormone (ACTH) into the blood, which can affect the adrenal gland. The effect of ACTH on the cortex of the adrenal gland is to elevate the release of cortisol, which will reduce inflammation and related processes [51, 52]. It seems that the alignment of the anti-inflammatory reflex and HPA axis can significantly inhibit inflammatory processes.

Therapeutic potential of using anti-inflammatory reflex

According to previous studies, using different methods to increase the stimulation of the vagus nerve or components related to the anti-inflammatory reflex can improve inflammatory disease [53, 30]. Clinical trials that activate the anti-inflammatory reflex include pharmacological and non-pharmacological interventions. Pharmacological treatments include acetylcholinesterase inhibitors, nicotine, and α 7nAChR. Non-pharmacological methods include invasive vagal nerve stimulation (VNS), non-invasive VNS, and pulsed ultrasound of the spleen [54]. Nicotine is a toxic agonist of all types of nicotinic receptors and has various side effects

[55]. The inhibition of acetylcholinesterase, an enzyme that is highly distributed in different cells, has many side effects that restrict its use. However, a suitable formulation of the α 7nAChR agonist has not yet been achieved in humans. Therefore, the use of drug therapy to activate inflammatory reflexes has several limitations. However, various studies have shown that VNS can reduce various inflammations [54]. For example, VNS has improved the inflammation of rheumatoid arthritis and lupus erythematosus [56, 57].

Heart rate variability; a natural phenomenon for evaluation of autonomic activity

The role of the autonomic nervous system in the anti-inflammatory reflex is clear. The status of the autonomic system can change depending on the activity of the anti-inflammatory reflex [58, 44]. Thus, when the anti-inflammatory reflex is activated, vagus nerve activity increases. One of the functions strongly affected by vagus nerve activity is heart rate [58]. Beat-to-beat heart rate variability is called HRV [59]. The autonomic nerves, especially the vagus nerve, are one of the main elements in creating heart rate variability [60]. It has been seen that in people with intensified inflammatory processes, the amount of HRV changes decreases [61]. During inflammation, the patient's HRV level fluctuates close to normal and the prognosis is considered positive. In addition, treatments that improve inflammation are associated with an increase in HRV, which indicates an increase in the activity of the vagus nerve [61, 62].

Although HRV is strongly influenced by the activity of the autonomic nerves, various factors can affect it, including hormones, drugs, and temperature [63]. At the same time, according to the spontaneous pacing of the heart, changes have been observed even in the isolated heart that has been inflamed, which shows the powerful effect of inflammation on all parts of the body [64, 65]. Therefore, by using HRV as an inexpensive, non-invasive, and accessible method, it is possible to evaluate the status of the autonomic system as the main component controlling the anti-inflammatory reflex.




Figure 1. Summary of anti-inflammatory reflex. The solid arrow indicates stimulation, whereas the dashed arrow indicates inhibition. Further details have been provided in the manuscript.

CONCLUSION

The anti-inflammatory reflexes are summarized in Figure 1. Inflammatory factors stimulate the afferent branches of the vagus nerve. The generated messages are processed in the central nervous system and the output is sent to macrophages and other white blood cells through the efferent pathway of the vagus nerve. Moreover, the HPA axis and adrenergic branch of the autonomic nerves in the spleen help control the inflammation (Figure 1). The activation of these responses to inflammation causes a reduction in the release of inflammatory factors, including cytokines. The generated negative feedback shows the interaction between the components of the nervous system and the immune cells, which together control the inflammatory processes. Inflammation is initially helpful for immune system function. However, with the increase in inflammation, positive feedback processes occur in which inflammatory factors enter the blood circulation and create a cascade of inflammatory reactions that are difficult for the body to control. Therefore, proper activity of the anti-inflammatory reflex leads to better management of inflammation along the HPA axis, which increases cortisol secretion.

Although the generality of this reflex has been identified, more research can help clarify other branches of its anti-inflammatory effects.

ACKNOWLEDGEMENT

The author is extremely grateful to Dr. Alireza Mani for his advice.

FUNDING

This research did not receive any specific grants from funding agencies in the public, commercial, or not-for-profit sectors.

DECLARATIONS

The author has no conflicts of interest to declare.

REFERENCES

- [1] Michael-Titus AT, Shortland P. The Nervous System: The Nervous System, E-Book. Elsevier Health Sciences; 2022.
- [2] Somjen G. Sensory coding in the mammalian nervous system. 2013.
- [3] Park KS. Nervous System. Humans and Electricity: Understanding Body Electricity and Applications. Springer; 2023. p. 27-51.

- [4] Ansar W, Ghosh S, Ansar W, Ghosh S. Inflammation and inflammatory diseases, markers, and mediators: Role of CRP in some inflammatory diseases. *Biology of C reactive protein in health and disease*. 2016;67-107.
- [5] Johnkennedy N, Mercy OC. Perspective of Inflammation and Inflammation Markers. *Journal La Medihealthico*. 2022;3(1):16-26.
- [6] Libby P. Inflammatory mechanisms: the molecular basis of inflammation and disease. *Nutrition reviews*. 2007;65(suppl_3):S140-S46.
- [7] Megari K. Quality of life in chronic disease patients. *Health psychology research*. 2013;1(3).
- [8] Gulati K, Guhathakurta S, Joshi J, Rai N, Ray A. Cytokines and their role in health and disease: a brief overview. *Moj Immunol*. 2016;4(2):00121.
- [9] Kany S, Vollrath JT, Relja B. Cytokines in inflammatory disease. *International journal of molecular sciences*. 2019;20(23):6008.
- [10] O'Shea JJ, Murray PJ. Cytokine signaling modules in inflammatory responses. *Immunity*. 2008;28(4):477-87.
- [11] De Jong HK, Van Der Poll T, Wiersinga WJ. The systemic pro-inflammatory response in sepsis. *Journal of innate immunity*. 2010;2(5):422-30.
- [12] Jaffer U, Wade R, Gourlay T. Cytokines in the systemic inflammatory response syndrome: a review. *HSR proceedings in intensive care & cardiovascular anesthesia*. 2010;2(3):161.
- [13] Fajgenbaum DC, June CH. Cytokine storm. *New England Journal of Medicine*. 2020;383(23):2255-73.
- [14] Sygitowicz G, Sitkiewicz D. Molecular mechanisms of organ damage in sepsis: an overview. *Brazilian Journal of Infectious Diseases*. 2021;24:552-60.
- [15] Tracey KJ. The inflammatory reflex. *Nature*. 2002;420(6917):853-59.
- [16] Hopkins SJ. Central nervous system recognition of peripheral inflammation: a neural, hormonal collaboration. *Acta Biomed*. 2007;78(Suppl 1):231-47.
- [17] Borovikova LV, Ivanova S, Zhang M, Yang H, Botchkina GI, Watkins LR, et al. Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature*. 2000;405(6785):458-62.
- [18] Pavlov VA, Tracey KJ. The cholinergic anti-inflammatory pathway. *Brain Behav Immun*. 2005 Nov;19(6):493-9.
- [19] Hoover DB. Cholinergic modulation of the immune system presents new approaches for treating inflammation. *Pharmacology & therapeutics*. 2017;179:1-16.
- [20] Bellocchi C, Carandina A, Montinaro B, Targetti E, Furlan L, Rodrigues GD, et al. The interplay between autonomic nervous system and inflammation across systemic autoimmune diseases. *International Journal of Molecular Sciences*. 2022;23(5):2449.
- [21] Bonaz B, Sinniger V, Pellissier S. Therapeutic potential of vagus nerve stimulation for inflammatory bowel diseases. *Frontiers in neuroscience*. 2021;15:650971.
- [22] Pavlov VA, Tracey KJ. The vagus nerve and the inflammatory reflex—linking immunity and metabolism. *Nature Reviews Endocrinology*. 2012;8(12):743-54.
- [23] Komegae EN, Farmer DGS, Brooks VL, McKinley MJ, McAllen RM, Martelli D. Vagal afferent activation suppresses systemic inflammation via the splanchnic anti-inflammatory pathway. *Brain, behavior, and immunity*. 2018;73:441-49.
- [24] Akira S. Toll-like receptors and innate immunity. *Advances in immunology*. 2001;78:1-56.
- [25] Karlsson H, Larsson P, Wold AE, Rudin A. Pattern of cytokine responses to gram-positive and gram-negative commensal bacteria is profoundly changed when monocytes differentiate into dendritic cells. *Scandinavian Journal of Immunology*. 2004;59(6):628-28.
- [26] Tietze K, Dalpke A, Morath S, Mutters R, Heeg K, Nonnenmacher C. Differences in innate immune responses upon stimulation with gram-positive and gram-negative bacteria. *Journal of periodontal research*. 2006;41(5):447-54.
- [27] Sternberg EM. Neural regulation of innate immunity: a coordinated nonspecific host response to pathogens. *Nature Reviews Immunology*. 2006;6(4):318-28.
- [28] Matsuda A, Jacob A, Wu R, Aziz M, Yang W-L, Matsutani T, et al. Novel therapeutic targets for sepsis: regulation of exaggerated inflammatory responses. *Journal of Nippon Medical School*. 2012;79(1):4-18.

- [29] Johnston G, Webster N. Cytokines and the immunomodulatory function of the vagus nerve. *British journal of anaesthesia*. 2009;102(4):453-62.
- [30] Bonaz B, Sinniger V, Pellissier S. Anti-inflammatory properties of the vagus nerve: potential therapeutic implications of vagus nerve stimulation. *The Journal of physiology*. 2016;594(20):5781-90.
- [31] Maier SF, Goehler LE, Fleshner M, Watkins LR. The role of the vagus nerve in cytokine-to-brain communication. *Annals of the New York Academy of Sciences*. 1998;840(1):289-300.
- [32] Maharjan A. Neuromodulation of the human autonomic nervous system using peripheral nerve stimulation techniques. University of Otago; 2022.
- [33] Garrett L, Trümbach D, Spielmann N, Wurst W, Fuchs H, Gailus-Durner V, et al. A rationale for considering heart/brain axis control in neuropsychiatric disease. *Mammalian Genome*. 2023;34(2):331-50.
- [34] Tanaka S, Hammond B, Rosin DL, Okusa MD. Neuroimmunomodulation of tissue injury and disease: an expanding view of the inflammatory reflex pathway. *Bioelectronic medicine*. 2019;5(1):13.
- [35] Pavlov VA, Tracey KJ. Neural circuitry and immunity. *Immunologic research*. 2015;63:38-57.
- [36] Matusik PS, Zhong C, Matusik PT, Alomar O, Stein PK. Neuroimaging studies of the neural correlates of heart rate variability: a systematic review. *Journal of clinical medicine*. 2023;12(3):1016.
- [37] Quadt L, Critchley H, Nagai Y. Cognition, emotion, and the central autonomic network. *Autonomic Neuroscience*. 2022;238:102948.
- [38] Rosas-Ballina M, Tracey K. Cholinergic control of inflammation. *Journal of internal medicine*. 2009;265(6):663-79.
- [39] Van Der Zanden EP, Boeckstaens GE, De Jonge WJ. The vagus nerve as a modulator of intestinal inflammation. *Neurogastroenterology & Motility*. 2009;21(1):6-17.
- [40] Metz CN, Pavlov VA. Vagus nerve cholinergic circuitry to the liver and the gastrointestinal tract in the neuroimmune communicome. *American Journal of Physiology-Gastrointestinal and Liver Physiology*. 2018;315(5):G651-G58.
- [41] Zdanowski R, Krzyżowska M, Ujazdowska D, Lewicka A, Lewicki S. Role of $\alpha 7$ nicotinic receptor in the immune system and intracellular signaling pathways. *Central European Journal of Immunology*. 2015;40(3):373-79.
- [42] Chao R, Tong Y-l, Li J-c, Lu Z-q, Yao Y-m. The protective effect of alpha 7 nicotinic acetylcholine receptor activation on critical illness and its mechanism. *International journal of biological sciences*. 2017;13(1):46.
- [43] Ren C, Li X-h, Wang S-b, Wang L-x, Dong N, Wu Y, et al. Activation of central alpha 7 nicotinic acetylcholine receptor reverses suppressed immune function of T lymphocytes and protects against sepsis lethality. *International Journal of Biological Sciences*. 2018;14(7):748.
- [44] Mazloom R, Eftekhari G, Rahimi-Balaei M, Khori V, Hajizadeh S, Dehpour AR, et al. The role of $\alpha 7$ nicotinic acetylcholine receptor in modulation of heart rate dynamics in endotoxemic rats. *PLoS One*. 2013;8(12):e82251.
- [45] Wu Y-j, Wang L, Ji C-f, Gu S-f, Yin Q, Zuo J. The role of $\alpha 7$ nAChR-mediated cholinergic anti-inflammatory pathway in immune cells. *Inflammation*. 2021;44(3):821-34.
- [46] Scanzano A, Cosentino M. Adrenergic regulation of innate immunity: a review. *Front Pharmacol*. 2015;6:171.
- [47] Pavlov VA. Collateral benefits of studying the vagus nerve in bioelectronic medicine. *Bioelectron Med*. 2019;5:5.
- [48] Durand M, Hagimont E, Louis H, Asfar P, Fripiat JP, Singer M, et al. The $\beta 1$ -Adrenergic Receptor Contributes to Sepsis-Induced Immunosuppression Through Modulation of Regulatory T-Cell Inhibitory Function. *Crit Care Med*. 2022 Sep 1;50(9):e707-e18.
- [49] McAllen RM, McKinley MJ, Martelli D. Reflex regulation of systemic inflammation by the autonomic nervous system. *Auton Neurosci*. 2022 Jan;237:102926.
- [50] Beishuizen A, Thijs LG. Endotoxin and the hypothalamo-pituitary-adrenal (HPA) axis. *J Endotoxin Res*. 2003;9(1):3-24.
- [51] DeMorrow S. Role of the Hypothalamic-Pituitary-Adrenal Axis in Health and Disease. *Int J Mol Sci*. 2018 Mar 26;19(4).
- [52] Miller WL. The Hypothalamic-Pituitary-Adrenal Axis: A Brief History. *Horm Res Paediatr*. 2018;89(4):212-23.

- [53] Boeckxstaens G. The clinical importance of the anti-inflammatory vagovagal reflex. *Handb Clin Neurol.* 2013;117:119-34.
- [54] Kelly MJ, Breathnach C, Tracey KJ, Donnelly SC. Manipulation of the inflammatory reflex as a therapeutic strategy. *Cell Rep Med.* 2022 Jul 19;3(7):100696.
- [55] Esmaeeli Dehaj H, Maleki Dehnavi S, Zahedi Nejad M, Akbarzadeh Kolahi S, Abdolghaffari AH, Khalili A, et al. The time interval between injection of nicotine and tremor initiation: a new index for evaluating nicotine efficacy in rodents. *Toxicol Mech Methods.* 2023 Dec 13:1-5.
- [56] Koopman FA, van Maanen MA, Vervoordeldonk MJ, Tak PP. Balancing the autonomic nervous system to reduce inflammation in rheumatoid arthritis. *J Intern Med.* 2017 Jul;282(1):64-75.
- [57] Aranow C, Atish-Fregoso Y, Lesser M, Mackay M, Anderson E, Chavan S, et al. Transcutaneous auricular vagus nerve stimulation reduces pain and fatigue in patients with systemic lupus erythematosus: a randomised, double-blind, sham-controlled pilot trial. *Ann Rheum Dis.* 2021 Feb;80(2):203-08.
- [58] Tracey KJ. Physiology and immunology of the cholinergic antiinflammatory pathway. *J Clin Invest.* 2007 Feb;117(2):289-96.
- [59] Camm AJ, Malik M, Bigger JT, Breithardt G, Cerutti S, Cohen RJ, et al. Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation.* 1996;93(5):1043-65.
- [60] Tiwari R, Kumar R, Malik S, Raj T, Kumar P. Analysis of Heart Rate Variability and Implication of Different Factors on Heart Rate Variability. *Curr Cardiol Rev.* 2021;17(5):e160721189770.
- [61] Scheff JD, Griffel B, Corbett SA, Calvano SE, Androulakis IP. On heart rate variability and autonomic activity in homeostasis and in systemic inflammation. *Math Biosci.* 2014 Jun;252:36-44.
- [62] Mol MBA, Strous MTA, van Osch FHM, Vogelaar FJ, Barten DG, Farchi M, et al. Heart-rate-variability (HRV), predicts outcomes in COVID-19. *PLoS One.* 2021;16(10):e0258841.
- [63] Billman GE. Heart rate variability—a historical perspective. *Frontiers in physiology.* 2011;2:86.
- [64] Gholami M, Mazaheri P, Mohamadi A, Dehpour T, Safari F, Hajizadeh S, et al. Endotoxemia is associated with partial uncoupling of cardiac pacemaker from cholinergic neural control in rats. *Shock.* 2012 Feb;37(2):219-27.
- [65] Mazloom R, Shirazi AH, Hajizadeh S, Dehpour AR, Mani AR. The effect of endotoxin on the controllability of cardiac rhythm in rats. *Physiol Meas.* 2014 Mar;35(3):339-49.